Nebraska Influenza & Other Respiratory Disease Surveillance Report, 2023-24 Influenza Season, Week 14 (DATA THROUGH WEEK ENDING 4/6). All data are preliminary and may change as more reports are received.

INFLUENZA WEEKLY SUMMARY

INFLUENZA LABORATORY SURVEILLANCE

Positive Influenza A & B Tests, Percent Positive, and Change from Last Week

Week Ending Date	Influenza A Positives	Change from Last Week	Influenza B Positives	Change from Last Week	Overall Percent Positive	% Change from Last Week
4/6/24	490	▼ 58	323	▼ 78	20.3%	▼1.8%
Grand Total	10,854		8,979			

Cumulative Influenza Positive Tests by Subtype and Age Group

	0-4	5-17	18-24	25-49	50-64	65+	Season Total
Flu A: H1	180	157	33	183	156	234	943
Flu A: H3	87	68	52	104	54	112	477
Flu B: Victoria	16	46	*	20	*	*	92

INFLUENZA-LIKE ILLNESS (ILI) OUTPATIENT SURVEILLANCE

Total ILI Visits Reported by the NE Outpatient ILI Surveillance Network (ILINet) and Change from Last Week

Week Ending Date	Total ILI Outpatient Visits	Change from Last Week
4/6/24	24	▼48
Grand Total	2,215	

SCHOOL ABSENTEEISM SURVEILLANCE

Percent of Students Absent due to any Illness, Number of Classroom/School Closures due to Illness, and Change from Last Week

Week Ending Date	% Absent (any illness)	% Change from Last Week	Classrooms Closed	Change from Last Week	Schools Closed	Change from Last Week
4/6/24	2.2%	▲ 0.2%	0	▼1	0	0

ILI EMERGENCY DEPARTMENT (ED) SURVEILLANCE

Total ILI ED Visits and Change from Last Week

Week Ending Date	Total ILI ED Visits	Change from Last Week
4/6/24	547	▼86
Grand Total	15,198	

ILI HOSPITALIZATION SURVEILLANCE

Total ILI Hospital Admissions and Change from Last Week

Week Ending Date	Total ILI Hospital Admissions	Change from Last Week
4/6/24	205	▼23
Grand Total	5,860	

LONG-TERM CARE FACILITY OUTBREAK SURVEILLANCE

27 influenza-associated outbreaks have been reported for the surveillance season

MORTALITY SURVEILLANCE

43 influenza-associated deaths have been reported for the surveillance season, including <6 pediatric deaths

National Influenza Summary: Please see http://www.cdc.gov/flu/weekly/

International Influenza Summary: Please see https://www.who.int/teams/global-influenza-programme/surveillance-and-monitoring/influenza-updates For information on the prevention of influenza, please see: http://www.cdc.gov/flu/protect/habits.htm

RESPIRATORY SYNCYTIAL VIRUS (RSV) WEEKLY SUMMARY

RSV LABORATORY SURVEILLANCE

Positive RSV Tests, Percent Positive, and Change from Last Week

Week Ending Date	RSV Positives	Change from Last Week	Percent Positive	% Change from Last Week
4/6/24	101	▼ 8	3.8%	▼0.1%
Grand Total	7 7/19			

RSV Percent Positive by Test Type and Percent Change from Last Week

	PCR		Anti	gen
4/6/24	3.6%	0.0%	4.9%	▼3.5%

LONG-TERM CARE FACILITY OUTBREAK SURVEILLANCE


16 RSV-associated outbreaks have been reported for the surveillance season


MORTALITY SURVEILLANCE


32 RSV-associated deaths have been reported for the surveillance season

(All data are preliminary and may change as more reports are received.)


INFLUENZA LABORATORY SURVEILLANCE








*The 2020 - 2021 influenza season was unusually low due much in part to the ongoing COVID-19 pandemic. As such, numbers for that season are substantially different than previous seasons and should be considered an anomaly.







Influenza Surveillance Data, Week 14 (Week Ending 4/6) (All data are preliminary and may change as more reports are received.)

INFLUENZA LABORATORY SURVEILLANCE, CONTINUED

Influenza A & B Positives by Test Type, by Week Ending Date, 2023-24







(All data are preliminary and may change as more reports are received.)


INFLUENZA LABORATORY SURVEILLANCE DEMOGRAPHICS


Influenza A & B Positives, by Age Group, by Week Ending Date, 2023-24



Influenza Positives by Patient Current Sex by Week Ending Date, 2023-24








Week Ending Date

Proportion of Influenza Positives by Patient Ethnicity, by Week Ending Date, 2023-24




Patient Race
Unknown
African American
Multiracial
American Indian/Alaskan Native
Native Hawaiian / Pacific Islander

Patient Ethnicity
Unknown Hispanic or Latino Non Hispanic or Latino

(All data are preliminary and may change as more reports are received.)

SCHOOL ABSENTEEISM SURVEILLANCE




Absenteeism Surveillance System Reporting Record over past 5 Weeks, 2023-24

(N=1,302 schools)

Week Ending Rep Date Er	orters	Number of Reports	Percent of Enrolled Reporting	Total Enrolled Reporters
3/9	0	839	83.3%	1,007
3/16	0	671	66.6%	1,007
3/23	-1	850	84.5%	1,006
3/30	0	840	83.5%	1,006
4/6	0	865	86.0%	1,006


Percentage of Students Absent due to any Illness and Number of Schools Closed due to Illness, by Week Ending Date, 2023-24



LONG-TERM CARE FACILITY OUTBREAK SURVEILLANCE

27 influenza-associated outbreaks have been reported for the surveillance season


INFLUENZA-LIKE ILLNESS (ILI) OUTPATIENT SURVEILLANCE



ILINet Sentinel Provider Reporting Record over the past 5 Weeks, 2023-24 (N=14, as of week ending 1/13)

	Number of Reports
3/9/24	12
3/16/24	11
3/23/24	11
3/30/24	10
4/6/24	8

 $Percentage \ of \ ILI\ Visits\ Reported\ by\ the\ Nebraska\ Outpatient\ ILI\ Surveillance\ Network\ (ILINet),\ by\ MMWR\ Week,\ 2020-2024$




(All data are preliminary and may change as more reports are received.)

INFLUENZA-LIKE ILLNESS (ILI) EMERGENCY DEPARTMENT (ED) SYNDROMIC SURVEILLANCE


Number of ILI Emergency Department (ED) Visits by Age Group, by Week Ending Date, 2023-24



Percentage of ILI Emergency Department Visits among all ED Visits by MMWR Week, 2020-2024




INFLUENZA-LIKE ILLNESS (ILI) HOSPITALIZATION SURVEILLANCE



ILI Hospital Reporting Record over the last 5 Weeks, 2023-24 (N=87 Hospitals)


Week Ending	n Date	Number of Hospitals Reporting	Percent of Hospitals Reporting
3/9/2	:4	67	76.1%
3/16/	/24	70	79.5%
3/23/	/24	64	72.7%
3/30/	/24	67	76.1%
4/6/2	<u>!</u> 4	57	64.8%

Number of ILI Admissions by MMWR Week, 2020-2024



MORTALITY SURVEILLANCE

43 influenza-associated deaths have been reported for the surveillance season, including <6 pediatric deaths Median Age: 70 years





RSV Surveillance Data, Week 14 (Week Ending 4/6)


All data are preliminary and may change as more reports are received.

RESPIRATORY SYNCYTIAL VIRUS (RSV) LABORATORY SURVEILLANCE

Number of Positive RSV Tests and Percent Positive by Week Ending Date, 2023-24







RSV Surveillance Data, Week 14 (Week Ending 4/6) (All data are preliminary and may change as more reports are received.)

RSV LABORATORY SURVEILLANCE, CONTINUED

RSV Percent Positive by Test Type, by Week Ending Date, 2023-24




Test Type Antigen PCR

Total RSV Tests by Test Type for past 10 Weeks

	2/3/24	2/10/24	2/17/24	2/24/24	3/2/24	3/9/24	3/16/24	3/23/24	3/30/24	4/6/24
Antigen	136	137	163	148	132	147	119	81	96	82
PCR	2,752	2,941	3,037	3,368	3,243	2,932	2,893	2,298	2,249	2,121


Week Ending Date




RSV Surveillance Data, Week 14 (Week Ending 4/6)

RSV LABORATORY SURVEILLANCE DEMOGRAPHICS






RSV Positives by Patient Current Sex, by Week Ending Date, 2023-24






Male



Proportion of RSV Positives by Patient Ethnicity, by Week Ending Date, 2023-24







RSV Surveillance Data, Week 14 (Week Ending 4/6)

(All data are preliminary and may change as more reports are received.)

OUTBREAK SURVEILLANCE

Age Group 0-4

16 RSV-associated outbreaks have been reported in long-term care facilities for the surveillance season

12/9/23


12/16/23

RSV EMERGENCY DEPARTMENT (ED) SYNDROMIC SURVEILLANCE

Week Ending Date

1/6/24


Percentage of RSV Emergency Department Visits among All ED Visits, by MMWR Week, 2020-2024



RSV INPATIENT SYNDROMIC SURVEILLANCE

5-24 25-49 50-64 65-

Percentage of RSV-Associated Inpatient Visits among All Inpatient Visits, by MMWR Week, 2020-2024



MORTALITY SURVEILLANCE

The state of the s

3/16/24


3/2/24

Other Respiratory Viruses Surveillance Data, Week 14 (Week Ending 4/6)

. (ΔII data are preliminary and may change as more reports are received.)

OTHER RESPIRATORY VIRUSES LABORATORY SURVEILLANCE

Cumulative PCR Positives by Respiratory Virus, 2023-24 season (starting 10/1/23)




Cumulative PCR Positives by Respiratory Virus and Virus Type, 2023-24


* 1 /		
Coronaviruses	Coronavirus 229E	176
	Coronavirus HKU1	252
	Coronavirus NL63	62
	Coronavirus OC43	527
	Coronavirus: Not Typed*	48
	Total	1,065
Human Metapneumovirus	Human Metapneumovirus	210
	Total	210
Parainfluenza Viruses	Parainfluenza 1	190
	Parainfluenza 2	99
	Parainfluenza 3	252
	Parainfluenza 4	122
	Parainfluenza: Not Typed*	15
	Total	678
Respiratory Adenovirus	Respiratory Adenovirus	848
	Total	848
Rhino/Enterovirus	Rhino/Enterovirus	3,600
	Total	3,600

*Not typed coronavirus and parainfluenza virus positives are NOT novel viruses. The specimens were simply were not typed out on their respective tests and both represent either known human seasonal coronaviruses (NOT SARS-CoV-2) or known seasonal human parainfluenza viruses respectively.

PCR Positives by Respiratory Virus and Virus Type, by Week Ending Date, 2023-24





PCR Positives by Respiratory Virus and Age Group, by Week Ending Date, 2023-24




Other Respiratory Viruses Surveillance Data, Week 14 (Week Ending 4/6)

OTHER RESPIRATORY VIRUSES LABORATORY SURVEILLANCE, CONTINUED






Respiratory Virus PCR Positives by Respiratory Season and MMWR Week, 2020-2024







About the Data

The Nebraska Influenza and other Respiratory Disease Surveillance System (NIRDSS) is a collaborative effort between DHHS and its many partners in the state including local health departments, public health and clinical laboratories, vital statistics offices, healthcare providers, clinics, and emergency departments.

Influenza surveillance allows us to determine when we first start to see influenza activity each year (the "first influenza case of the season"), and provides an indicator of the progression of the influenza season as well as prevalence of disease in the community, which assists healthcare providers in diagnosing patients with influenza-like illness (ILI). ILI is defined as any patient with clinically diagnosed influenza or any patient with fever $\geq 100^{\circ}F$ ($\geq 37.8^{\circ}C$), oral or equivalent, AND cough and/or sore throat. The case definition no longer includes "without a known cause other than influenza". Surveillance additionally identifies what strains of influenza are circulating in any given year, and thus determines whether the current vaccine protects against the circulating strain. By incorporating multiple data sources, we are able to communicate a more complete picture of influenza activity.

For information about Morbidity and Mortality Weekly Report (MMWR) weeks, please see: https://ndc.services.cdc.gov/wp-content/uploads/MMWR_week_overview.pdf

For the 2022-2023 MMWR Week Calendar, please see:

https://ndc.services.cdc.gov/wp-content/uploads/MMWR-Week-Log-2022-2023.pdf

NOTE: Data values of 1-5 are supressed throughout this report for patient confidentiality purposes. These values are denoted with '*' or '<6'.

<u>Laboratory Surveillance</u>

The Nebraska Laboratory Influenza Surveillance Program consists of hospital-based laboratories that submit testing data, either by weekly survey or daily electronic laboratory report (ELR). These laboratories perform rapid antigen or PCR testing for influenza and Respiratory Syncytial Virus (RSV). The Nebraska Public Health Laboratory provides further characterization of a subset of influenza isolates to determine the subtype of influenza A viruses and the lineage of influenza B viruses. Influenza A subtypes are determined by proteins, hemagglutinin (H) and neuraminidase (N), found on the outside of the virus. For the purpose of this report, influenza A subtypes are categorized into two groups, H1 and H3, as these two subtypes most commonly circulate during influenza season. Influenza B lineages are classified into one of two lineages: Yamagata and Victoria. The age, patient current sex, race, ethnicity, and test type data figures in the laboratory surveillance section utilizes ELR data only. The age, gender, race, and ethnicity data is obtained directly from lab reports; data missing from lab reports or specifically listed as unknown in the lab report are combined as "Unknown" in this report. All other data figures in this section utilize ELR data and laboratory data received via survey from Nebraska labs who do not participate in ELR.

Many influenza and RSV disease cases are never reported. Most people with influenza or RSV do not see a doctor about their illness. Many of those who do seek care are not tested, and only a portion of test results that are obtained are reported to DHHS. DHHS receives laboratory reports from facilities participating in automated electronic laboratory reporting. We do not receive reports on all positive tests. Because some providers actively test for influenza and others do not, relying solely on case counts for influenza could result in an incomplete assessment of influenza community activity.

When testing for respiratory illnesses, there are two tests most commonly used in practices. The first of the two is an antigen test, which is most common between the two. Antigen tests are inexpensive tests that generally take only 15-30 minutes to return with results. Antigen tests try to identify specific proteins on the surface of the virus. The other type of test is a polymerase chain reaction (PCR) test. This test tries to identify specific genetic material for the virus. PCR tests take longer to produce results compared to antigen tests, but it is considered the gold standard for testing because it is a lot more sensitive than the antigen test.

Note on RSV Percent Positive: An antigen test positivity of 10% and a polymerase chain reaction (PCR) test positivity of 3% are accepted threshold levels for determining when RSV activity is considered to be at an epidemic level. The healthcare community monitors these test positivity thresholds, and when they are surpassed it indicates RSV activity is increasing throughout the population. These signals give healthcare providers more insight to know when to begin recommending monoclonal antibody therapy (i.e. Palivizumab and Nirsevimab) to infants to protect them from severe illness due to RSV. More information on these therapies can be found here: https://www.cdc.gov/vaccines/vpd/rsv/immunization-information-statement.html

All data presented for the "OTHER RESPIRATORY VIRUSES LABORATORY SURVEILLANCE" is obtained from our ELR data. This data only includes PCR tests, and a majority of these PCR tests are PCR respiratory virus panels. This data is limited to the number of laboratories who participate in ELR. Furthermore, historical data may be limited due to a fewer number of laboratories participating in ELR compared to more recent years, making it more difficult to compare data from recent years to years further in the past.

About the Data, Continued

School Absenteeism Surveillance

The School Absenteeism Surveillance System captures data on the total expected enrollment at Nebraska schools, the number of total absences, and the number of absences due to specific illnesses, like influenza and COVID-19. This surveillance system is also used to alert local health departments if absenteeism is above 10% which could indicate an outbreak situation. This system is designed to encourage communication between schools and local health departments and to promote the accessibility of Nebraska's public health system if schools need assistance, for example, with potential disease outbreaks. This data is analyzed and reported for the current surveillance week so potential outbreak situations can be identified and responded to in a timely manner.

A school closure is when an entire school is closed (all students and staff are sent home or a switch to virtual learning). A classroom closure is if the school is open for most students, but, due to an outbreak cluster in a particular classroom, only the students / staff in that classroom are absent.

For more information on preventing outbreaks in schools, visit: https://www.cdc.gov/flu/school/guidance.htm

Long-Term Care Facility Outbreak Surveillance

Reporting of influenza outbreaks in long-term care facilities (LTCF), schools and other congregate settings is required by rules and regulations.

173 NAC 1 1-004.01B Clusters, Outbreaks, or Unusual Events, Including Possible Bioterroristic Attacks: Clusters, outbreaks, or epidemics of any health problem, infectious or other, including food poisoning, healthcare-associated outbreaks or clusters, influenza, or possible bioterroristic attack; increased disease incidence beyond expectations; unexplained deaths possibly due to unidentified infectious causes; and any unusual disease or manifestations of illness must be reported immediately.

<u>Definition of respiratory outbreak (not COVID-19):</u>

A sudden increase in acute febrile respiratory illness* over the normal background rate (e.g., 2 or more cases of acute respiratory illness occurring within 72 hours of each other)

*Acute febrile respiratory illness is defined as fever > 100°F AND one or more respiratory symptoms (runny nose, sore throat, laryngitis, or cough). However, please note that elderly patients with influenza may not develop a fever.

Nebraska Outpatient ILI Surveillance (ILINet)

Voluntary reporting by a statewide network of sentinel clinicians of the number of patients presenting with influenza-like illness (ILI) and the total number of patient visits by age group each week.

Emergency Department and Inpatient Syndromic Surveillance

The NE Syndromic Surveillance System monitors influenza-like and RSV-associated illness data received by 71/85 Nebraska emergency departments and 64/88 Nebraska inpatient facilities. Syndromic surveillance is the real-time (or near real-time) collection of patient visits to a Nebraska emergency department where discharge diagnoses and/or chief complaint include influenza and RSV-associated illness.

ILI Hospitalization Surveillance

Voluntary reporting by hospital infection preventionists of the number of hospitalizations with a diagnosis of ILI and the total number of admissions by age group each surveillance week.

Mortality Surveillance

Pediatric deaths associated with influenza are required to be reported. Influenza-associated deaths in adults are not reportable. RSV-associated deaths are not reportable of any age. Voluntary reporting to public health of deaths in adults is encouraged to help determine the severity of the current circulating virus.