



United States  
**Consumer Product Safety Commission**

# **Non-Fire Carbon Monoxide Deaths Associated with the Use of Consumer Products 2019 Annual Estimates**

March 2023

John Topping

Directorate for Epidemiology

Division of Hazard Analysis

U.S. Consumer Product Safety

4330 East West Highway

Bethesda, MD 20814

*This report was prepared by the CPSC staff.  
It has not been reviewed or approved by,  
and may not necessarily reflect the views of,  
the Commission.*

# Table of Contents

<b>Executive Summary</b> .....	<b>4</b>
<b>Introduction</b> .....	<b>6</b>
<b>National Estimates of Non-Fire CO Poisoning Deaths Associated with Consumer Products</b> .....	<b>7</b>
Table 1: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Associated Fuel-Burning Consumer Product, 2009–2019.....	11
Table 1: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Associated Fuel-Burning Consumer Product, 2009–2019 (continued) .....	12
Table 1: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Associated Fuel-Burning Consumer Product, 2009–2019 (continued) .....	13
Table 2: Estimated Non-Fire Carbon Monoxide Poisoning Deaths Associated with Consumer Products Organized by Fuel Type, 2009–2019.....	15
Table 2: Estimated Non-Fire Carbon Monoxide Poisoning Deaths Associated with Consumer Products Organized by Fuel Type, 2009–2019 (continued) .....	16
Table 2: Estimated Non-Fire Carbon Monoxide Poisoning Deaths Associated with Consumer Products Organized by Fuel Type, 2009–2019 (continued) .....	17
Table 3: Estimated Non-Fire Carbon Monoxide Poisoning Deaths Associated with Engine-Driven Tools, 2009–2019.....	19
Figure 1: Comparison of Trends in Consumer Product-Related Carbon Monoxide Deaths, 2009 to 2019 .....	21
Table 4: Number of Carbon Monoxide Poisoning Incidents Reported to CPSC by Number of Deaths per Incident, 2009–2019.....	23
Table 5: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Location of Exposure, 2009–2019 .....	25
Table 6: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Month and Year of the Fatality, 2009–2019.....	27
Figure 2: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Month of Death, 2009–2019 .....	28
<b>Victim Demographics from Non-Fire Carbon Monoxide Poisoning Deaths Associated with the Use of Consumer Products</b> .....	<b>29</b>
Table 7: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Age of Victim, 2009–2019 .....	30
Table 8: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Gender of Victim, 2009–2019 .....	31

Table 9: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Race/Ethnicity, 2009–2019 .....	33
Table 10: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Race/Ethnicity, 2009–2019 .....	35
Table 11: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Race/Ethnicity, 2009–2019 .....	36
<b>Appendix A: Methodology .....</b>	<b>37</b>
<b>Appendix B: National Estimates and Mortality Rates of Consumer Product-Related CO Poisoning Deaths, 1980 to 2018.....</b>	<b>44</b>
<b>Appendix C: Chi-Squared Test Results.....</b>	<b>52</b>
<b>Appendix D: Regional Definitions.....</b>	<b>53</b>
<b>References .....</b>	<b>54</b>

# Executive Summary

This report provides information about the estimated number of unintentional non-fire deaths attributed to carbon monoxide (CO) poisoning that were associated with the use of consumer products in 2019, along with companion statistics since 2009. Because the U.S. Consumer Product Safety Commission (CPSC) has not received reports of deaths from every state for 2018 and 2019, the estimates for those years may change in subsequent reports.

Some of the key findings<sup>1</sup> in this report are:

## For 2019:

- CPSC has records from 168 incidents resulting in an estimated 250 unintentional non-fire CO poisoning deaths associated with the use of consumer products under the CPSC's jurisdiction.
- Fourteen percent of the 168 incidents involved multiple deaths, including two incidents where four people died, and another two incidents where three people died.
- Engine-Driven Tools (EDTs) were associated with the largest percentage of non-fire CO poisoning deaths, more than any other category. This category includes generators, the single product under CPSC's jurisdiction that is associated with the most CO deaths. Just under half (118 or 47%) of the estimated 250 deaths associated with consumer products involved an EDT. One hundred of the 118 estimated EDT-associated deaths involved generators,
- Heating Systems were associated with the second largest percentage of non-fire CO poisoning deaths. An estimated 69 deaths (28%) were associated with some type of heating appliance. Seventy-five percent of the estimated 250 CO deaths in 2019 resulted from CO exposure in a home location. Within incidents coded as home locations, a few deaths resulted from CO exposure in an external structure at a residence (e.g., detached garage), a non-fixed location domicile used as a home (e.g., camper trailers), a structure not designed for habitation but used as a home (e.g., metal shed), as well as tents, or temporary shelters.
- In 2019, males constituted 77 percent of CO poisoning victims.

## For 2017-2019:

- The estimated annual average from 2017 to 2019 was 216 deaths.
- Most CO deaths occurred in the colder months of the year, with more than half of the deaths occurring during the four cold months of November, December, January, and February.
- Adults 45 years and older comprised an annual average of more than 66 percent of all non-fire, consumer product-related CO deaths, which was disproportionately higher for this age group than their representation in the U.S. population. Conversely, children younger than 15 years of age accounted for a disproportionately lower annual average of less than 4 percent of the yearly CO poisoning deaths.

---

<sup>1</sup> Note that the estimates for individual categories may not sum to that of the broader category due to rounding effects.

- In the period 2017-2019, some statistical evidence demonstrates that the proportion of deaths by race/ethnicity differs from the proportions of race/ethnicity in the U.S. population. The proportion of Hispanic victims (irrespective of race) is significantly lower than the proportion of Hispanic Americans in the U.S. population; in contrast, the proportion of White victims was significantly greater than their percentage in the U.S. population.

**For 2009-2019:**

- Staff found evidence of a statistically significant upward trend in non-fire CO deaths for the 11-year period from 2009 to 2019. The estimated number of consumer product-related CO deaths in 2019 is greater than any other year in this report; in fact, the estimate in this category is greater than any estimate since the changeover from ICD-9 to ICD-10 in 1999. The estimate has increased now for the seventh straight year.
- Since 2009, portable generators alone have been associated with an estimated 765 non-fire CO poisoning deaths, accounting for 40 percent of all CO deaths related to consumer products under CPSC's jurisdiction.

## Introduction

Carbon monoxide (CO) is a colorless, odorless, and poisonous gas that results from the incomplete combustion of fuels, such as natural or liquefied petroleum (LP) gas, gasoline, oil, wood, coal, and other fuels. The health effects related to CO depend upon its concentration in blood, which, in turn, depends upon its concentration in air, an individual's duration of exposure, and an individual's general health. Carbon monoxide combines with the body's hemoglobin (Hb) with an affinity about 250 times that of oxygen, forming carboxyhemoglobin (COHb) and interfering with oxygen uptake, delivery, and use by the cells. Generally, no perceptible health effects or symptoms in healthy individuals occur at COHb levels below 10 percent. Symptoms associated with blood levels at or above 10 percent COHb include headache, fatigue, nausea, and cognitive impairment. Loss of consciousness, coma, and death can occur at COHb levels greater than 20 percent; but for healthy adults, CO deaths typically require levels above 50 percent COHb.<sup>2</sup> Staff notes that during exposure to rapidly rising, high CO levels (as can result with exposure to exhaust from gasoline-powered, engine-driven tools), sudden extreme hypoxia can result in rapid incapacitation and loss of consciousness, which prevent exposed individuals from leaving the CO environment.

Some symptoms of CO poisoning may mimic common illnesses, such as influenza or colds. Thus, a possibility of initial misdiagnosis by physicians and victims exists (Long and Saltzman, 1995). Frequently, patients are unaware of exposures, and health care providers may not always consider CO poisoning as a cause of such nonspecific symptoms. COHb formation is reversible, as are some clinical symptoms of CO poisoning. However, some delayed neurological effects that develop after severe poisonings, especially those involving prolonged unconsciousness, may not be reversible. Prompt medical attention is important to reduce the risk of permanent damage.

Any fuel-burning appliance can be a potential source of fatal or hazardous CO levels. Fuels, such as natural and LP gas, kerosene, oil, coal, and wood, can produce large amounts of CO when insufficient oxygen is available for combustion. Consumer products that burn kerosene, oil, coal, or wood (such as wood stoves, oil boilers, and kerosene heaters) often produce an irritating smoke that can sometimes alert the victim to a potentially hazardous situation. EDTs powered by gasoline engines produce large amounts of CO, even in locations where sufficient oxygen is available for combustion. However, EDTs may not emit an irritating exhaust smoke. Other fuels, such as charcoal briquettes and pressed wood-chip logs produce relatively smokeless fires, even at times of inefficient combustion. In these cases, victims receive no obvious sensory warning that can alert victims to a potentially hazardous situation. Another hazard scenario is present when gas appliances are not vented properly or are malfunctioning. Natural and LP gas burn more efficiently and cleanly, compared to other forms of fuel. However, in circumstances of poor maintenance, inadequate ventilation, or faulty exhaust pathways, natural and LP gas appliances may emit potentially lethal amounts of CO without any irritating fumes. Again, many victims may be unaware of a potential problem.

# National Estimates of Non-Fire CO Poisoning Deaths Associated with Consumer Products

The national estimates presented in this report are based on death certificate records obtained from all 50 states, the District of Columbia, New York City, and some U.S. territories, directly augmented by information collected in CPSC's In-Depth Investigations (IDIs), and to a lesser extent, news articles, and medical examiners' reports contained in the CPSC Injury or Potential Injury Incident (IPII) database. Death certificate data from some states can lag for months, or even years, and may not be available in time for use in this report.

The estimates of consumer product-related CO poisoning deaths presented in this report are based on reporting up to September 1, 2022. The National Center for Health Statistics (NCHS) has records of every death certificate filed in the United States and its territories. Before 2017, there was evidence that CPSC records contained a large portion of the records reported to NCHS. For the years 2008 through 2015, CPSC records contain approximately 82 percent of all the fatal CO poisoning deaths that occurred in the United States as reported to NCHS. However, in 2016, and to a slightly lesser extent in 2017, there appears to have been an anomaly with the method used by Texas in assigning ICD-10 codes used in this analysis, in particular, the Y17 code (see Appendix A for details on the methodology used to determine estimates). The estimates presented here are based on the number of deaths for which CPSC has records, scaled to the NCHS totals, to adjust for missing records. Appendix A of this report describes the detailed process used to generate the national estimates presented in this report.

It also should be noted that, due to extended reporting delays from a small proportion of U.S. states, a potentially significant portion of death certificates are missing from the 2018 and 2019 data. Although most states apparently completed reporting through 2019, there seems to be not even a single death certificate captured in CPSC databases from the state of Washington for the year 2018 or 2019. This seems to reflect delays considerably beyond most other states that have not reported additional 2018 or prior deaths since July 2021, when data were already consolidated for last year's version of this report. Consequently, estimates for years 2018 and earlier remain unchanged since last year. There also appear to be anomalies with reporting from Wisconsin and Texas, highly suggestive of incomplete capture of relevant death certificates from those states for 2018 and 2019. Some adjustments were made using historical patterns to account for anomalies in previous years 2016 and 2017 for the state of Texas, when there was no comparable anomaly for the state of Wisconsin. For the years 2018 and 2019, there are not yet sufficient data to support special adjustments for any of these three states, and therefore, the estimates should be considered incomplete for those most recent 2 years. See additional discussion on this topic in Appendix A.

During 2019, an estimated 250 non-fire CO poisoning deaths were associated with the use of a consumer product under the jurisdiction of the CPSC. This report does not include CO poisoning deaths involving products outside CPSC's jurisdiction, such as incidents where the CO gas resulted from a fire, or solely from a motor vehicle, or directly work related; and the report also does not include deaths that were suicides or otherwise intentional in nature. Over the prior 10 years, the annual average was 166 estimated non-fire CO deaths from consumer

products. Please note that the estimates and findings for the 11 years covered in this report, include three incidents (one each in 2013, 2015, and 2016), where the exhaust from a motor vehicle engine may have contributed to the victim's CO poisoning death, in addition to a consumer product. Additionally, in another included 2016 incident, a farm tractor may have contributed to a CO fatality, along with an unspecified lawn mower that was running in a residential storage shed. Utility vehicles and ATVs are considered consumer products (not motor vehicles); and therefore, CO from their exhaust is considered relevant, regardless. For example, in 2019, four such CO deaths due to exhaust (3 from utility vehicles and 1 from an ATV) are classified as Off-Highway Vehicles (OHVs), as a subcategory under Engine-Driven-Tools.

Although multiple factors may contribute to a CO poisoning fatality, the source of CO is virtually always a fuel-burning product. The following factors can cause or contribute to a fuel-burning product producing dangerous levels of CO: poor product design, product failure or malfunction; improper service or maintenance; improper venting of exhaust products; consumer misuse; inadequate ventilation of the room in which the product is located. CPSC staff produces the CO estimates associated with consumer products, to identify and monitor product groups involved in these fatal CO scenarios. Within the individual product-specific CPSC projects, additional analysis assesses whether improvements are warranted in the areas of product design, ventilation safeguards, or user information and education.

The annual CO estimates for the years 2009 through 2019, are presented in two formats: by product category (Table 1), and by product within fuel type (Table 2). The data are presented as an average of the most recent 3-year period (2017 through 2019), followed by yearly estimates for each of the 11 years covered by this report. As noted, collection of death certificates from some states is incomplete for 2018 and 2019. Accordingly, although reporting for those years is complete from most states through 2019, estimates for those years may change, if additional data become available, in particular, from non-reporting states. Therefore, data for 2018 and 2019 are reported using italic font in the tables, to signify reporting is incomplete.

Because the numbers presented in this document represent national estimates of unintentional, non-fire deaths attributed to CO poisoning associated with the use of consumer products, the generator and other EDT death estimates would not be expected to match the observed fatality counts presented in this report or in the CPSC report, "Fatal and Nonfatal Incidents Associated with Non-Fire Carbon Monoxide from Engine-Driven Generators and Other Engine-Driven Tools, 2011–2021," published in August 2022.

## **By Product Category**

Table 1 shows the estimated average annual number of CO poisoning deaths associated with various consumer products for 2017 to 2019, as well as the annual estimated CO deaths for the individual years from 2009 through 2019. The annual average for 2017 through 2019 is estimated to be 216 (with a standard error of approximately 9.8). Appendix B contains a graph and the data point values for the annual estimates of CO poisoning deaths associated with a consumer product under CPSC's jurisdiction for 1980 through 2019.

The estimate for Heating Systems, which historically account for a large percentage of the deaths, is further broken down into heating system subcategories within various fuel types. Fatality estimates for the Engine-Driven Tools category were distributed between generators and other engine-driven tools. The consumer product-related estimate and estimate-by-product distribution were derived using the methodology described in Appendix A.

In 2019, products in the Heating Systems category were associated with an estimated 69 deaths (28% of the total 250 CO poisoning deaths from consumer products). Of the 69, the majority (86% or 59 deaths) were known to have involved gas heating systems or devices. Natural gas heating was associated with an estimated 17 deaths (25% of all heating system-related deaths). LP gas heating was associated with an estimated 40 deaths (58% of heating system-related deaths); and unspecified gas heating was associated with an estimated two deaths (3% of heating system-related deaths).

Staff notes that several other fuel-burning devices, not specifically designed for heating purposes, were known or suspected of having been used for heating an enclosed space where a victim died of CO poisoning. Such devices included charcoal/charcoal grills (an estimated 14 deaths) and gas ranges (3 deaths).

Of the estimated 17 deaths associated with natural gas (NG) heating, the majority (76% or 13 deaths) involved installed freestanding furnaces. The remaining four involved an NG wall furnace. At least half of the estimated 40 deaths in 2019 that were associated with LP gas heating systems involved unvented portable propane heaters, not including an additional 10 (25%) that involved an unidentified LP heating device. The unvented, portable propane heaters were fueled by a propane tank and were not a component of an installed heating system. The portable LP heaters are intended as camping heaters or heaters for other temporary spaces and use disposable, refillable, or exchangeable propane tanks.

There were also an estimated two deaths associated with coal-burning heating devices: one from a coal furnace, and one from a coal-burning stove.

Additionally, in 2019:

- An estimated four deaths were associated with some type of heating system where the fuel was unknown, one product was known to be a furnace, and three products simply were identified as heating systems.
- An estimated 14 CO deaths (6% of the 250 total estimated deaths) were associated with charcoal or charcoal grills. As noted, most of these were either known to have been used, or were suspected of being used for heating purposes, often in temporary spaces, like inside a vehicle.
- An estimated 18 deaths (7%) were associated with residential water heaters, where an estimated 10 were propane-fueled, three were natural gas water heaters, and five were fueled by an unspecified gas.
- An estimated 16 deaths were associated with multiple appliances (6% of the total estimated deaths). The multiple-products category includes all incidents where multiple fuel-burning products were used simultaneously, such that a single source of the CO could not be determined.

In recent years, the Engine-Driven Tools category, which includes generators, lawn mowers, leaf blowers, tillers, power washers, and snow blowers, among other EDTs, has been associated with more CO deaths than any other category. Nearly half of the estimated average number of CO deaths in the three most recent years (2017 through 2019) were associated with engine-driven tools (107 of 216, not including multiple-product incidents). Over the 11 years covered in this report, the total number of estimated CO deaths associated with engine-driven tools (893) exceeds the estimates for heating systems (562). Estimated generator-related CO deaths alone exceed those for heating systems over these 11 years (765 generator-related deaths, versus 562 heating system-related deaths). When a single CO-producing product is involved, generator-related deaths comprise most engine-driven, tools-related CO deaths, accounting for 86 percent of all engine-driven tools-related deaths over the entire 11 years covered by this report.

The availability of detailed information regarding the condition of products associated with CO deaths varies widely. Information collected often describes conditions indicative of compromised vent systems, flue passageways, and chimneys for furnaces, boilers, and other heating systems. Vent systems include the portion of piping that either connects the flue outlet of the appliance and exhausts air to the outside through a ceiling or sidewall or connects to a chimney. According to the information available, some products had vents that became detached or were installed/maintained improperly. Vents were also sometimes blocked by soot caused by inefficient combustion, which, in turn, may have been caused by several factors, such as leaky or clogged burners, an over-firing condition, or inadequate combustion air.

Other reported furnace-related conditions included compromised heat exchangers or filter doors/covers that were removed or not sealed. Some products were old and apparently not well maintained. Other incidents mentioned a backdraft condition, large amounts of debris in the chimney, and the use of a product that was later prohibited by the utility company and designated not to be turned on until repaired.

**Table 1: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Associated Fuel-Burning Consumer Product, 2009–2019**

Consumer Product	2017–2019+		Annual Estimates										
	Average Estimate	Average Percent	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018+	2019+
<b>Total</b>	<b>216</b>	<b>100%</b>	<b>148</b>	<b>159</b>	<b>163</b>	<b>137</b>	<b>146</b>	<b>164</b>	<b>172</b>	<b>178</b>	<b>188</b>	<b>210</b>	<b>250</b>
<b>Heating Systems</b>	<b>58</b>	<b>27%</b>	<b>41</b>	<b>58</b>	<b>49</b>	<b>46</b>	<b>43</b>	<b>64</b>	<b>37</b>	<b>50</b>	<b>41</b>	<b>64</b>	<b>69</b>
<b>Furnaces (incl. Boilers)</b>	<b>22</b>	<b>10%</b>	<b>16</b>	<b>30</b>	<b>22</b>	<b>27</b>	<b>21</b>	<b>24</b>	<b>20</b>	<b>34</b>	<b>17</b>	<b>27</b>	<b>22</b>
Coal	1	1%	*	*	1	*	*	*	*	*	2	1	1
Liquid Petroleum (LP) Gas	2	1%	1	7	*	4	1	11	3	3	*	*	5
Natural Gas	13	6%	10	15	6	15	5	6	6	15	6	21	13
Oil	1	<1%	3	1	2	*	5	1	3	2	1	1	*
Unspecified Gas	2	1%	1	4	10	4	10	6	8	11	4	*	1
Unspecified Fuel	3	1%	1	2	2	5	*	*	1	2	4	4	1
<b>Portable Heaters</b>	<b>22</b>	<b>10%</b>	<b>8</b>	<b>19</b>	<b>13</b>	<b>11</b>	<b>12</b>	<b>18</b>	<b>11</b>	<b>11</b>	<b>19</b>	<b>26</b>	<b>21</b>
Kerosene	1	<1%	*	1	2	1	*	2	1	4	2	*	1
Liquid Petroleum (LP) Gas	21	10%	8	18	11	10	12	14	10	6	16	26	20
Natural Gas	*	*	*	*	*	*	*	1	*	*	*	*	*
Unspecified Gas	*	*	*	*	*	*	*	*	*	1	*	*	*
Unspecified Fuel	<1	<1%	*	*	*	*	*	*	*	*	1	*	*
<b>Wall/Floor Furnaces</b>	<b>3</b>	<b>1%</b>	<b>6</b>	<b>5</b>	<b>1</b>	<b>*</b>	<b>*</b>	<b>5</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>1</b>	<b>5</b>
Liquid Petroleum (LP) Gas	<1	<1%	5	1	*	*	*	*	*	1	*	*	1
Natural Gas	2	1%	1	2	*	*	*	2	1	*	*	1	4
Unspecified Gas	<1	<1%	*	*	*	*	*	2	*	*	1	*	*
Unspecified Fuel	<1	<1%	*	1	1	*	*	*	*	*	1	*	*
<b>Room/Space Heaters</b>	<b>6</b>	<b>3%</b>	<b>9</b>	<b>1</b>	<b>5</b>	<b>5</b>	<b>9</b>	<b>8</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>6</b>	<b>10</b>
Coal	1	<1%	*	*	2	*	1	1	*	*	*	1	1
Liquid Petroleum (LP) Gas	2	1%	5	1	1	4	3	7	*	*	*	*	7
Natural Gas	*	*	2	*	*	*	2	*	*	*	*	*	*
Wood	2	1%	2	*	1	*	2	*	*	*	1	4	1
Unspecified Gas	<1	<1%	*	*	1	*	*	*	*	1	*	1	*
Unspecified Fuel	<1	<1%	*	*	*	1	*	*	1	*	*	*	1
<b>Unspecified Heater/System</b>	<b>5</b>	<b>2%</b>	<b>2</b>	<b>4</b>	<b>8</b>	<b>2</b>	<b>1</b>	<b>9</b>	<b>3</b>	<b>3</b>	<b>1</b>	<b>4</b>	<b>10</b>
Liquid Petroleum (LP) Gas	3	1%	*	1	3	1	*	8	1	*	*	2	7
Natural Gas	*	*	*	*	1	*	*	*	*	*	*	*	*
Unspecified Gas	<1	<1%	1	1	1	1	*	*	*	*	*	*	1
Unspecified Fuel	2	1%	1	1	2	*	1	1	1	3	1	2	3
<b>Charcoal Grills, Charcoal</b>	<b>11</b>	<b>5%</b>	<b>7</b>	<b>17</b>	<b>10</b>	<b>6</b>	<b>11</b>	<b>7</b>	<b>11</b>	<b>7</b>	<b>10</b>	<b>10</b>	<b>14</b>

**Table 1: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Associated Fuel-Burning Consumer Product, 2009–2019 (continued)**

Consumer Product	2017–2019+		Annual Estimates										
	Average Estimate	Average Percent	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018+	2019+
<b>Engine-Driven Tools</b>	<b>107</b>	<b>50%</b>	<b>76</b>	<b>56</b>	<b>73</b>	<b>64</b>	<b>68</b>	<b>62</b>	<b>92</b>	<b>80</b>	<b>104</b>	<b>100</b>	<b>118</b>
Generators – Gasoline	78	36%	64	40	64	57	55	53	84	61	89	72	74
Generators – LP	*	*	*	2	*	*	*	*	*	7	*	*	*
Generators – Unspecified Fuel	14	7%	*	*	*	*	1	1	*	*	6	11	26
Other Engine-Driven Tools	15	7%	12	14	10	6	13	8	8	12	10	16	18
<b>Ranges or Ovens</b>	<b>8</b>	<b>4%</b>	<b>4</b>	<b>5</b>	<b>8</b>	<b>4</b>	<b>10</b>	<b>*</b>	<b>5</b>	<b>7</b>	<b>12</b>	<b>9</b>	<b>3</b>
Liquid Petroleum (LP) Gas	1	<1%	*	1	1	1	1	*	3	*	1	1	*
Natural Gas	2	1%	2	2	3	*	2	*	3	6	*	2	3
Unspecified Gas	5	2%	2	1	3	2	2	*	*	*	11	5	*
Unspecified Fuel	*	*	*	*	*	*	5	*	*	1	*	*	*
<b>Water Heaters</b>	<b>9</b>	<b>4%</b>	<b>5</b>	<b>2</b>	<b>8</b>	<b>5</b>	<b>2</b>	<b>5</b>	<b>9</b>	<b>6</b>	<b>4</b>	<b>5</b>	<b>18</b>
Liquid Petroleum (LP) Gas	4	2%	2	*	1	*	1	1	*	*	*	2	10
Natural Gas	3	1%	1	2	4	*	*	*	*	1	4	1	7
Oil	*	*	*	*	*	*	*	*	*	*	*	*	*
Unspecified Gas	2	1%	1	*	1	2	*	2	8	4	*	1	1
Unspecified Fuel	*	*	1	*	1	2	1	1	1	*	*	*	*
<b>Pool Heaters</b>	<b>1</b>	<b>&lt;1%</b>	<b>*</b>	<b>1</b>	<b>1</b>	<b>*</b>	<b>3</b>	<b>2</b>	<b>*</b>	<b>2</b>	<b>4</b>	<b>*</b>	<b>*</b>
Liquid Petroleum (LP) Gas	*	*	*	1	*	*	*	*	*	*	*	*	*
Natural Gas	1	<1%	*	*	*	*	3	1	*	*	4	*	*
Unspecified Gas	*	*	*	*	1	*	*	*	*	2	*	*	*
Unspecified Fuel	*	*	*	*	*	*	*	1	*	*	*	*	*
<b>Lanterns</b>	<b>*</b>	<b>*</b>	<b>1</b>	<b>*</b>	<b>2</b>	<b>2</b>	<b>*</b>	<b>5</b>	<b>5</b>	<b>1</b>	<b>*</b>	<b>*</b>	<b>*</b>
Liquid Petroleum (LP) Gas	*	*	1	*	1	2	*	4	5	1	*	*	*
Kerosene	*	*	*	*	*	*	*	1	*	*	*	*	*
Unspecified Fuel	*	*	*	*	1	*	*	*	*	*	*	*	*
<b>Grills, Camp Stoves</b>	<b>5</b>	<b>2%</b>	<b>*</b>	<b>*</b>	<b>2</b>	<b>*</b>	<b>1</b>	<b>6</b>	<b>4</b>	<b>3</b>	<b>6</b>	<b>4</b>	<b>4</b>
Liquid Petroleum (LP) Gas	3	1%	*	*	2	*	*	2	4	3	6	1	1
Coal	<1	<1%	*	*	*	*	*	*	*	*	*	1	*
Wood	1	<1%	*	*	*	*	*	1	*	*	*	*	3
Unspecified Fuel	<1	<1%	*	*	*	*	1	2	*	*	*	1	*

Note: Use of a natural gas water heater not as the product was intended with Liquid Petroleum (LP) gas (instead of natural gas) is associated with an estimated 4 deaths in 2019. Those deaths are classified based on *product type* under “Water Heaters -- natural gas”, even though LP gas was used.

**Table 1: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Associated Fuel-Burning Consumer Product, 2009–2019 (continued)**

Consumer Product	2017–2019+		Annual Estimates										
	Average Estimate	Average Percent	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018+	2019+
<b>Other Products</b>	<b>5</b>	<b>2%</b>	<b>2</b>	<b>5</b>	<b>3</b>	<b>2</b>	<b>2</b>	<b>7</b>	<b>1</b>	<b>2</b>	<b>1</b>	<b>7</b>	<b>8</b>
Chimney – Unspecified Fuel	<i>1</i>	<i>&lt;1%</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>1</i>	<i>1</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>3</i>
Fire Pit – Wood	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>1</i>	<i>*</i>	<i>*</i>	<i>*</i>
Fireplace – Coal	<i>*</i>	<i>*</i>	<i>*</i>	<i>1</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>
Other Products – LP Gas	<i>2</i>	<i>1%</i>	<i>1</i>	<i>1</i>	<i>2</i>	<i>*</i>	<i>1</i>	<i>4</i>	<i>1</i>	<i>*</i>	<i>1</i>	<i>4</i>	<i>*</i>
Other Products – Natural Gas	<i>*</i>	<i>*</i>	<i>1</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>
Other Products – Unspecified Fuel	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>1</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>
Unidentified Product	<i>2</i>	<i>1%</i>	<i>*</i>	<i>2</i>	<i>1</i>	<i>1</i>	<i>*</i>	<i>1</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>1</i>	<i>5</i>
Unidentified Product – LP Gas	<i>1</i>	<i>&lt;1%</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>*</i>	<i>1</i>	<i>*</i>	<i>1</i>	<i>*</i>	<i>2</i>	<i>*</i>
<b>Multiple Products</b>	<b>11</b>	<b>5%</b>	<b>11</b>	<b>15</b>	<b>8</b>	<b>5</b>	<b>5</b>	<b>7</b>	<b>9</b>	<b>19</b>	<b>6</b>	<b>12</b>	<b>16</b>

+ Data collection for 2018 and 2019 is only partially complete, and data are shown in italics. Italicized estimates may change in the future if more reports of deaths are received.

\* No reports received by CPSC staff.

Source: U.S. Consumer Product Safety Commission/EPHA.

CPSC Death Certificate File, CPSC Injury or Potential Injury Incident File, CPSC In-Depth Investigation File, National Center for Health Statistics Mortality File, 2009–2019.

Note: Reported annual estimates and estimated averages and percentages may not add to subtotals or totals due to rounding.

## By Fuel Type

Table 2 (beginning on page 18) organizes the estimates, by product, within fuel type. The three major fuel types include: *Gas-Fueled Products* (natural gas and liquid petroleum—LP including propane and butane—gas); *Solid-Fueled Products* (charcoal, coal, and wood); and *Liquid-Fueled Products* (gasoline, kerosene, and oil). Of these fuel types, *Gas-Fueled Products* were associated with 89 of the 250 (36%) estimated CO deaths in 2019. *Liquid-Fueled Products* were associated with an estimated 97 (39%) deaths; and *Solid-Fueled Products* were associated with an estimated 21(8%) deaths in the same period. There were also 40 fatalities (16%), where the fuel type of the device could not be identified.

In the *Gas-Fueled Products* category in 2019, an estimated 68 of the 89 gas-fueled appliance deaths (76%) were associated with heating systems or heaters, including furnaces, portable heaters, and room or space heaters. Additionally, all eight of the *Multiple Gas-Fueled Products* fatalities were associated with a heating-related product and another product.

All but four of the estimated 97 liquid-fueled, appliance-related deaths in 2019, were associated with engine-driven tools (e.g., generators, lawn mowers/garden tractors). An estimated 74 deaths were associated with gasoline-fueled generators.

In 2019, an estimated 21 deaths fit within the *Solid-Fueled Products* category. Fourteen of these were associated with charcoal or charcoal grills.

**Table 2: Estimated Non-Fire Carbon Monoxide Poisoning Deaths Associated with Consumer Products Organized by Fuel Type, 2009–2019**

Consumer Product	2017–2019 <sup>+</sup>		Annual Estimates										
	Average Estimate	Average Percent	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018 <sup>+</sup>	2019 <sup>+</sup>
<b>Total</b>	<b>216</b>	<b>100%</b>	<b>148</b>	<b>159</b>	<b>163</b>	<b>137</b>	<b>146</b>	<b>164</b>	<b>172</b>	<b>178</b>	<b>188</b>	<b>210</b>	<b>250</b>
<b>Gas-Fueled Products</b>	<b>76</b>	<b>35%</b>	<b>53</b>	<b>70</b>	<b>58</b>	<b>51</b>	<b>45</b>	<b>78</b>	<b>53</b>	<b>69</b>	<b>60</b>	<b>78</b>	<b>89</b>
<b>Natural Gas</b>	<b>21</b>	<b>10%</b>	<b>17</b>	<b>23</b>	<b>15</b>	15	13	11	10	21	13	23	26
Furnace (incl. Boilers)*	12	6%	10	15	6	15	5	6	6	15	6	18	13
Pool Heater	1	1%	*	*	*	*	3	1	*	*	4	*	*
Portable Heater	*	*	*	*	*	*	*	1	*	*	*	*	*
Range/Oven	2	1%	2	2	3	*	2	*	3	6	*	2	3
Room/Space Heater	*	*	2	*	*	*	2	*	*	*	*	*	*
Wall/Floor Furnace	2	1%	1	2	*	*	*	2	1	*	*	1	4
Water Heater	4	2%	1	2	4	*	*	*	*	1	4	1	7
Unspecified Heater	*	*	*	*	1	*	*	*	*	*	*	*	*
Other Appliance	*	*	1	*	*	*	*	*	*	*	*	*	*
<b>Liquid Petroleum (LP) Gas</b>	<b>39</b>	<b>18%</b>	<b>23</b>	<b>35</b>	<b>23</b>	<b>22</b>	<b>20</b>	<b>52</b>	<b>27</b>	<b>22</b>	<b>24</b>	<b>41</b>	<b>51</b>
Furnace (incl. Boilers)	2	1%	1	7	*	4	1	11	3	3	*	2	5
Generator	<1	<1%	*	2	*	*	*	*	*	7	*	1	*
Grill/Camp Stove	3	1%	*	*	2	*	*	2	4	3	6	1	1
Lantern	*	*	1	*	1	2	*	4	5	1	*	*	*
Other Products/Unknown	1	1%	1	*	*	*	*	2	1	1	*	4	*
Pool Heater	*	*	*	1	*	*	*	*	*	*	*	*	*
Portable Heater	20	9%	8	18	11	10	12	14	10	6	16	24	20
Range/Oven	1	0%	*	1	1	1	1	*	3	*	1	1	*
Refrigerator	1	0%	*	1	2	*	1	2	*	*	1	2	*
Room/Space Heater	2	1%	5	1	1	4	3	7	*	*	*	*	7
Unspecified Heater/System	3	1%	*	1	3	1	*	8	1	*	*	2	7
Wall/Floor Furnace	<1	<1%	5	1	*	*	*	*	*	1	*	*	1
Water Heater	4	2%	2	*	1	*	1	1	*	*	*	2	10
<b>Unspecified Gas</b>	<b>9</b>	<b>4%</b>	<b>5</b>	<b>6</b>	<b>17</b>	<b>10</b>	<b>13</b>	<b>11</b>	<b>15</b>	<b>20</b>	<b>16</b>	<b>7</b>	<b>4</b>
Furnace (incl. Boilers)	2	1%	1	4	10	4	10	6	8	11	4	*	1
Pool Heater	*	*	*	*	1	*	*	*	*	2	*	*	*
Portable Heater	*	*	*	*	*	*	*	*	*	1	*	*	*
Range/Oven	5	2%	2	1	3	2	2	*	*	*	11	5	*
Room/Space Heater	<1	<1%	*	*	1	*	*	*	*	1	*	1	*
Fireplace	*	*	*	*	*	*	*	*	*	*	*	*	*
Wall/Floor Furnace	<1	<1%	*	*	*	*	*	2	*	*	1	*	*
Water Heater	1	0%	1	*	1	2	*	2	8	4	*	1	1
Unspecified Heater	<1	<1%	1	1	1	1	*	*	*	*	*	*	1

**Table 2: Estimated Non-Fire Carbon Monoxide Poisoning Deaths Associated with Consumer Products Organized by Fuel Type, 2009–2019 (continued)**

Consumer Product	2017–2019*		Annual Estimates										
	Average Estimate	Average Percent	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018*	2019*
<b>Multiple Gas-Fueled Products</b>	7	3%	8	6	3	4	*	5	1	6	6	6	8
<b>Liquid-Fueled Products</b>	96	44%	81	60	79	65	73	67	96	83	102	89	97
<b>Gasoline-Fueled</b>	92	42%	77	53	73	64	67	61	92	73	98	86	91
Generator	78	36%	64	40	64	57	55	53	84	61	88	72	74
Other Engine-Driven Tools	14	6%	12	14	10	6	13	8	8	12	10	15	17
<b>Kerosene-Fueled</b>	1	<1%	*	1	2	1	*	4	1	4	2	*	1
Portable Heater	1	<1%	*	1	2	1	*	2	1	4	2	*	1
Lantern	*	*	*	*	*	*	*	1	*	*	*	*	*
<b>Oil-Fueled</b>	1	<1%	3	1	2	*	5	1	3	2	1	1	*
Furnace (incl. Boilers)	1	<1%	3	1	2	*	5	1	3	2	1	1	*
Water Heater	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Diesel-Fueled</b>	*	*	*	*	*	*	*	*	*	*	*	*	*
Water Heater	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Multiple Liquid-Fueled Products</b>	2	1%	1	5	1	*	1	1	*	3	*	1	5
<b>Solid-Fueled Products</b>	17	8%	9	18	14	5	14	9	11	8	13	17	21
<b>Charcoal-Fueled</b>	11	5%	7	17	10	5	11	7	11	7	10	10	14
Charcoal/Charcoal Grills	11	5%	7	17	10	5	11	7	11	7	10	10	14
<b>Coal-Fueled</b>	3	1%	*	1	3	*	1	1	*	*	2	4	3
Furnace (incl. Boilers)	1	<1%	*	*	1	*	*	*	*	*	2	1	1
Room/Space Heater	1	<1%	*	*	2	*	1	1	*	*	*	1	1
Coal Grill/Coal	<1	<1%	*	*	*	*	*	*	*	*	*	1	*
Chimney/Fireplace	*	*	*	1	*	*	*	*	*	*	*	*	*
<b>Wood-Fueled</b>	3	1%	2	*	1	*	2	1	*	1	1	4	4
Fire Pit	*	*	*	*	*	*	*	*	*	1	*	*	*
Grill/Stove	1	<1%	*	*	*	*	*	1	*	*	*	*	3
Room/Space Heater	2	1%	2	*	1	*	2	*	*	*	1	4	1

**Table 2: Estimated Non-Fire Carbon Monoxide Poisoning Deaths Associated with Consumer Products Organized by Fuel Type, 2009–2019 (continued)**

Consumer Product	2017–2019+		Annual Estimates										
	Average Estimate	Average Percent	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018+	2019+
<b>Unspecified Fuel Products</b>	<b>25</b>	<b>11%</b>	<b>3</b>	<b>7</b>	<b>9</b>	<b>11</b>	<b>10</b>	<b>8</b>	<b>5</b>	<b>7</b>	<b>13</b>	<b>21</b>	<b>40</b>
Chimney	1	<1%	*	*	*	1	1	*	*	*	*	*	3
Furnace (incl. Boilers)	3	1%	1	2	2	5	*	*	1	2	4	4	1
Generator	14	7%	*	*	*	*	1	1	*	*	6	11	26
Grill/Camp Stove	<1	<1%	*	*	*	*	1	2	*	*	*	1	*
Lantern	*	*	*	*	1	*	*	*	*	*	*	*	*
Pool Heater	*	*	*	*	*	*	*	1	*	*	*	*	*
Portable Heater	<1	<1%	*	*	*	*	*	*	*	*	1	*	*
Range/Oven	*	*	*	*	*	*	5	*	*	1	*	*	*
Room/Space Heater	<1	<1%	*	*	*	1	*	*	1	*	*	*	1
Unspecified Heater	2	1%	1	1	2	*	1	1	1	3	1	2	2
Wall/Floor Furnace	<1	<1%	*	1	1	*	*	*	*	*	1	*	*
Unidentified Product	2	1%	*	2	1	1	*	1	*	*	*	1	4
OEDT	<1	<1%	*	*	*	*	*	*	*	*	*	1	1
Water Heater	*	*	1	*	1	2	1	1	1	*	*	*	*
<b>Multiple Product - Different Fuels</b>	<b>2</b>	<b>1%</b>	<b>2</b>	<b>4</b>	<b>3</b>	<b>4</b>	<b>3</b>	<b>1</b>	<b>8</b>	<b>10</b>	<b>*</b>	<b>5</b>	<b>2</b>
Gas & Liquid	2	1%	1	1	2	2	3	1	6	8	*	5	2
Gas & Solid	*	*	1	*	*	1	*	*	*	1	*	*	*
Liquid & Solid	*	*	*	2	1	*	*	*	*	*	*	*	*
Liquid & Unspecified	*	*	*	*	*	*	*	*	1	1	*	*	*

+ Data collection for 2018 and 2019 is only partially complete. Italicized estimates may change in the future if more reports of deaths are received.

\* No reports received by CPSC staff.

Source: U.S. Consumer Product Safety Commission/EPHA.

CPSC Death Certificate File, CPSC Injury or Potential Injury Incident File, CPSC In-Depth Investigation File, National Center for Health Statistics Mortality File, 2009–2019.

Note: Use of a natural gas water heater not as the product was intended with Liquid Petroleum (instead of natural gas) is associated with an estimated 4 deaths in 2019. Those deaths are classified based on product type under “natural gas” [water heater], even though LP gas was used.

## Engine-Driven Tools

Table 3 shows a breakdown of the fatality estimates for the 11-year period from 2009 through 2019, in the *Engine-Driven Tools* category. During 2019, engine-driven tools were associated with an estimated 123 carbon monoxide poisoning deaths (49% of the 250 total consumer product-related CO death estimate). In the 3 most recent years, EDTs comprised 51 percent of all consumer-product-related CO fatalities (estimated annual average of 111 out of 216 – when including several multiple product deaths associated with a generator and/or EDT). Of these EDT fatalities, generators dominated, with an estimated annual average of 93 out of 111.

Lawnmowers were associated with slightly less than half of the deaths listed in the *Other Engine-Driven Tools* category for the 11-year period (58 of 127 total fatalities). There were six other deaths associated with a lawnmower and another product in this period. There was an estimated average of six lawnmower-related CO deaths per year from 2017 to 2019 (18 deaths, excluding multiproduct deaths). There were multiple fatalities for six other sub-categories over the 2017 to 2019 period: power washers (an estimated 6 fatalities), OHV (6), leaf blowers (2), snow blowers (2), and unspecified EDTs (2).

**Table 3: Estimated Non-Fire Carbon Monoxide Poisoning Deaths Associated with Engine-Driven Tools, 2009–2019**

Engine-Driven Tools	2017-2019+		Annual Estimate										
	Average Estimate	Average Percentage	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018+	2019+
<b>Total</b>	<i>111</i>	<i>100%</i>	<b>78</b>	<b>61</b>	<b>78</b>	<b>66</b>	<b>73</b>	<b>64</b>	<b>97</b>	<b>92</b>	<b>104</b>	<b>106</b>	<b>123</b>
<b>Generators</b>	<i>93</i>	<i>84%</i>	<b>64</b>	<b>42</b>	<b>64</b>	<b>57</b>	<b>56</b>	<b>54</b>	<b>84</b>	<b>67</b>	<b>95</b>	<b>84</b>	<b>100</b>
Gasoline-fueled	<i>78</i>	<i>70%</i>	64	40	64	57	55	53	84	61	88	72	74
LP-fueled	<i>&lt;1</i>	<i>&lt;1%</i>	*	<b>2</b>	*	*	*	*	*	7	*	<b>1</b>	*
Unspecified Fuel	<i>14</i>	<i>13%</i>	*	*	*	*	1	1	*	*	6	11	26
<b>Other Engine-Driven Tools (OEDTs)</b>	<i>15</i>	<i>14%</i>	<b>12</b>	<b>14</b>	<b>10</b>	<b>6</b>	<b>13</b>	<b>8</b>	<b>8</b>	<b>12</b>	<b>10</b>	<b>16</b>	<b>18</b>
<b>Lawn Mowers</b>	<i>6</i>	<i>5%</i>	<b>6</b>	<b>7</b>	<b>3</b>	<b>4</b>	<b>7</b>	<b>2</b>	<b>4</b>	<b>7</b>	<b>5</b>	<b>9</b>	<b>4</b>
Riding Mowers	<i>3</i>	<i>3%</i>	6	5	3	2	6	1	3	7	4	5	*
Unspecified Mowers	<i>3</i>	<i>3%</i>	*	2	*	1	1	1	1	*	1	4	4
Paint Sprayer	<i>*</i>	<i>*</i>	*	*	1	*	*	*	*	*	*	*	*
Power Washer	<i>2</i>	<i>2%</i>	1	*	2	*	*	2	*	1	1	1	4
Snow Blower/Thrower	<i>1</i>	<i>1%</i>	3	1	1	*	2	1	1	1	*	1	1
OHV (e.g., ATV or UTV)	<i>2</i>	<i>2%</i>	*	4	2	1	1	1	*	*	1	*	5
Water Pump	<i>&lt;1</i>	<i>&lt;1%</i>	*	1	*	*	1	*	*	*	*	*	1
Welder	<i>*</i>	<i>*</i>	*	*	*	*	*	*	3	1	*	*	*
Tiller	<i>&lt;1</i>	<i>&lt;1%</i>	1	*	*	*	*	*	*	*	*	1	*
Leaf Blower	<i>1</i>	<i>1%</i>	*	*	*	*	*	*	*	1	*	2	*
Go-Cart	<i>*</i>	<i>*</i>	1	*	*	*	*	*	*	*	*	*	*
Antique Tractor	<i>&lt;1</i>	<i>&lt;1%</i>	*	*	*	*	*	*	*	*	1	*	*
Small Engine	<i>*</i>	<i>*</i>	*	*	*	*	*	*	*	1	*	*	*
Snowmobile	<i>&lt;1</i>	<i>&lt;1%</i>	*	*	*	*	*	*	*	*	*	*	1
Stump Grinder	<i>&lt;1</i>	<i>&lt;1%</i>	*	*	*	*	1	*	*	*	1	*	*
Wood Splitter	<i>*</i>	<i>*</i>	*	*	*	1	*	1	*	*	*	*	*
Unspecified EDT	<i>1</i>	<i>1%</i>	*	*	*	*	*	*	*	*	*	1	1
<b>Multiple Product: Engine-Driven Tools Involved</b>	<i>4</i>	<i>4%</i>	<b>2</b>	<b>6</b>	<b>4</b>	<b>2</b>	<b>5</b>	<b>2</b>	<b>5</b>	<b>12</b>	<b>*</b>	<b>6</b>	<b>5</b>
Generator + OEDT	<i>&lt;1</i>	<i>&lt;1%</i>	*	*	*	*	*	*	*	1	*	*	1
Generator + other Product	<i>3</i>	<i>3%</i>	2	6	3	2	3	2	4	10	*	6	4
Multiple OEDT	<i>*</i>	<i>*</i>	*	*	1	*	*	*	*	1	*	*	*
OEDT + other product	<i>*</i>	<i>*</i>	*	*	*	*	1	*	1	*	*	*	*

+ Data collection for 2018 and 2019 is only partially complete, and data are shown in italics. Italicized estimates may change in the future if more reports of deaths are received.

\* No reports received by CPSC staff.

Source: U.S. Consumer Product Safety Commission/EPHA.


CPSC Death Certificate File, CPSC Injury or Potential Injury Incident File, CPSC In-Depth Investigation File, National Center for Health Statistics Mortality File, 2009–2019.

Note: Reported annual estimates and estimated averages and percentages may not add to subtotals or totals due to rounding.

## Comparison of Trends

Figure 1 provides a graphic representation of the CO fatality trends related to: (1) all consumer products; (2) engine-driven tools; and (3) non-engine-driven tool products. A regression analysis of the estimated number of all non-fire, consumer product-related CO poisoning deaths from 2009 to 2019, indicates evidence of a statistically significant trend (p-value = 0.0015). The estimated CO fatalities from consumer products has now risen for the seventh straight year. In 2018, the estimated number of CO fatalities had risen above 200 deaths for the first time since before the changeover from ICD-9 to ICD-10 in 1999, and the estimate increased to 250 in the year 2019. Part of this is likely due to an uptick in CO fatalities associated with engine-driven tools.

**Figure 1: Comparison of Trends in Consumer Product-Related Carbon Monoxide Deaths, 2009 to 2019**



Source: U.S. Consumer Product Safety Commission/EPHA.  
CPSC Death Certificate File, CPSC Injury or Potential Injury Incident File, CPSC In-Depth Investigation File, 2009–2019.

## Number of Deaths per Incident Reported to CPSC

Table 4 presents a summary of the incident data broken down by the number of deaths per incident. Staff notes that this table does **not** provide estimates; the numbers presented are counts observed in the CPSC databases. Therefore, the counts presented in Table 4 should not be expected to add up to the estimated deaths in other tables. Table 4 shows that in 2019, 145 of the 168 fatal CO incidents (85% of fatal CO incidents reported to the CPSC) involved a single death. Table 4 accounts for only the fatally injured victims in each CO poisoning incident. It is not uncommon for CO incidents involving one or more deaths to also result in one or more nonfatal CO poisoning injuries. However, the breakdown of these injuries was not quantified for analysis in this death-focused report.

Occasionally, even though CPSC records indicate that there was more than one fatality in a specific incident, not all the deaths are used in the estimation process. Deaths for which CPSC does not have a death certificate are not used in the analyses, because the scaling estimation process accounts for missing records. Also, if an additional fatality is recorded as work related, that fatality is not counted in the estimation process, because work-related deaths are out of scope for this report. However, both scenarios are included in Table 4 to highlight the danger of multiple deaths in CO poisoning cases.

Death certificates do not include information about other deaths for the same incident. The number of deaths for a particular incident is based primarily on CPSC In-Depth Investigation (IDI) records. Some additional multiple-fatality incidents were identified by matching the incident date of death and location of death to the death certificate, while others were identified from news articles contained in the CPSC Injury or Potential Injury Incident (IPII) database. Over the 11-year period covered by this report, CPSC records indicate that 18 percent of the incidents resulted in multiple deaths. Nineteen incidents resulted in four or more CO deaths, including an incident in 2015, where eight people died, and another incident in 2016, in which six people died.

**Table 4: Number of Carbon Monoxide Poisoning Incidents Reported to CPSC by Number of Deaths per Incident, 2009–2019**

Number of Deaths Reported in Incident	2017–2019 <sup>+</sup>		Annual Incidents										
	Annual Average	Average Percent	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018 <sup>+</sup>	2019 <sup>+</sup>
<b>All Incidents</b>	<b><i>148</i></b>	<b><i>100%</i></b>	<b><i>117</i></b>	<b><i>116</i></b>	<b><i>120</i></b>	<b><i>90</i></b>	<b><i>106</i></b>	<b><i>110</i></b>	<b><i>104</i></b>	<b><i>130</i></b>	<b><i>131</i></b>	<b><i>145</i></b>	<b><i>168</i></b>
1	<i>125</i>	<i>85%</i>	93	100	95	74	84	86	83	110	112	<i>119</i>	<i>145</i>
2	<i>19</i>	<i>13%</i>	19	14	22	14	21	21	15	16	15	<i>23</i>	<i>19</i>
3	<i>2</i>	<i>2%</i>	4	1	1	1	*	1	2	1	3	<i>2</i>	<i>2</i>
4	<i>1</i>	<i>1%</i>	1	1	1	*	1	1	3	2	1	<i>1</i>	<i>2</i>
5	*	*	*	*	1	1	*	1	*	*	*	*	*
6	*	*	*	*	*	*	*	*	*	1	*	*	*
7	*	*	*	*	*	*	*	*	*	*	*	*	*
8	*	*	*	*	*	*	*	*	1	*	*	*	*

+ Data collection for 2018 and 2019 is only partially complete, and data are shown in italics. Italicized counts may change in the future if more reports of deaths are received.

Note: Percentages do not add to 100% due to rounding.

Numbers presented here are counts based on records available to CPSC staff. These do not represent national estimates and should not be expected to match estimates presented elsewhere in this document.

Source: U.S. Consumer Product Safety Commission/EPHA

CPSC Death Certificate File, CPSC Injury or Potential Injury Incident File, CPSC In-Depth Investigation File.

## By Location of Exposure

Table 5 shows that in 2019, an estimated 187 CO poisoning deaths resulted from exposure to CO in home locations, including an estimated 15 deaths from CO in detached structures at residential locations (*i.e.*, sheds, detached garages); and another 18 from CO in structures not intended originally as a permanent residence (*i.e.*, camper trailers, sea-land shipping containers). From 2017 to 2019, an estimated annual average of 164 deaths (76% of the annual average estimate for all CO deaths) resulted from exposure to CO in home locations. In 2019, an estimated 21 deaths resulted from CO in temporary shelters, such as campers, cabins, and trailers used for shelter. For 2017 to 2019, an annual average of 22 deaths (10%) resulted from CO in temporary shelters. Deaths due to CO exposures in temporary shelters were most associated with heating sources or generators.

A small percentage of the CO poisoning deaths resulted from CO in vehicles (such as passenger vans, trucks, automobiles, or boats), where a consumer product was the CO-producing product in use. In 2019, there were an estimated 19 CO deaths in this category. For the 3-year period 2017 to 2019, an annual average of 17 deaths (8%) resulted from CO in vehicles. Vehicle location incidents in this 3-year period usually involved a generator, LP heater, grill, or the burning of charcoal inside the vehicle.

**Table 5: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Location of Exposure, 2009–2019**

Location of Exposure	2017–2019 <sup>+</sup>		Annual Estimate										
	Average Estimate	Average Percent	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018 <sup>+</sup>	2019 <sup>+</sup>
<b>Total</b>	<b>216</b>	<b>100%</b>	<b>148</b>	<b>159</b>	<b>163</b>	<b>137</b>	<b>146</b>	<b>164</b>	<b>172</b>	<b>178</b>	<b>188</b>	<b>210</b>	<b>250</b>
Home <sup>1</sup>	135	62%	109	125	122	107	104	100	113	135	142	108	154
Home – External Structure <sup>2</sup>	12	6%	7	5	10	5	13	15	14	11	5	17	15
Home – But Not House <sup>3</sup>	17	8%	1	5	5	1	3	12	4	3	11	23	18
Temporary Shelter	22	10%	18	17	15	21	16	21	24	19	13	33	21
Vehicles (including boats)	17	8%	12	6	9	*	7	6	12	4	10	22	19
Outdoors	6	3%	*	*	*	*	*	*	*	1	*	1	11
Other	9	4%	*	1	1	*	2	8	5	1	7	4	17
Unknown	3	1%	*	*	1	2	*	1	*	1	*	2	4

+ Data collection for 2018 and 2019 is only partially complete, and data are shown in italics. Italicized estimates may change in the future if more reports of deaths are received.

\* No reports received by CPSC staff.

Note: Percentages do not add to 100% due to rounding.

1 Traditional home (e.g., detached house, townhouse, apartment, mobile home)

2 External structure at residential locations (e.g., detached garage, shed)

3 Non-fixed structure or structure not originally designed for permanent occupation (e.g., camper trailer, van, converted sea-land shipping container).

Source: U.S. Consumer Product Safety Commission/EPHA.

CPSC Death Certificate File, CPSC In-Depth Investigation File, CPSC Injury or Potential Injury Incident File, National Center for Health Statistics Mortality File, 2009–2019.

## **By Time of Year**

CPSC data indicate that there were more CO deaths attributable to incidents that occurred in the cold months than in the warm months. This is most likely because of the use of furnaces and portable heaters in the cold months. Additionally, generators are often used in the cold months because of power outages due to snow and ice storms. Table 6 shows the annual estimated CO deaths categorized by month of death. In 2019, an estimated 137 of the 250 estimated CO deaths (55%) were attributable to deaths that occurred during the four cold months of November, December, January, and February. Over the 11 years covered by this report, the average percentage of deaths occurring in the four colder months is 55 percent. In 2019, an estimated 73 deaths (29%) are attributable to incidents that occurred during the transition months of March, April, September, and October. This is only slightly lower than the 11-year average of 30 percent for the same four months. And in the warmer months of May, June, July, and August, an estimated 40 CO deaths (16%) occurred.

**Table 6: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Month and Year of the Fatality, 2009–2019**

Month of Death	2017–2019 <sup>+</sup>		Annual Estimate										
	Average Estimate	Average Percent	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018 <sup>+</sup>	2019 <sup>+</sup>
<b>Total</b>	<b>216</b>	<b>100%</b>	<b>148</b>	<b>159</b>	<b>163</b>	<b>137</b>	<b>146</b>	<b>164</b>	<b>172</b>	<b>178</b>	<b>188</b>	<b>210</b>	<b>250</b>
<b>Cold Months</b>	<b>112</b>	<b>52%</b>	<b>85</b>	<b>109</b>	<b>85</b>	<b>75</b>	<b>82</b>	<b>83</b>	<b>82</b>	<b>109</b>	<b>80</b>	<b>120</b>	<b>137</b>
November	24	11%	12	18	34	26	16	20	10	32	17	27	27
December	25	11%	20	38	20	25	28	20	23	19	21	29	24
January	35	16%	29	38	24	10	22	26	24	28	27	38	40
February	29	14%	24	15	8	14	16	17	24	29	16	26	46
<b>Transition Months</b>	<b>65</b>	<b>30%</b>	<b>41</b>	<b>33</b>	<b>55</b>	<b>46</b>	<b>43</b>	<b>44</b>	<b>62</b>	<b>49</b>	<b>65</b>	<b>56</b>	<b>73</b>
March	16	7%	12	22	9	6	12	10	19	12	13	12	22
April	16	7%	8	6	11	14	6	14	28	13	14	15	18
September	16	7%	4	2	13	6	5	6	11	7	23	20	5
October	17	8%	17	2	23	20	21	14	4	17	14	10	27
<b>Warm Months</b>	<b>39</b>	<b>18%</b>	<b>21</b>	<b>17</b>	<b>23</b>	<b>16</b>	<b>21</b>	<b>37</b>	<b>29</b>	<b>20</b>	<b>42</b>	<b>34</b>	<b>40</b>
May	10	5%	5	8	9	2	4	17	4	5	9	11	10
June	14	6%	10	5	2	5	6	4	9	3	16	14	12
July	7	3%	4	2	4	7	7	13	11	6	10	1	10
August	8	4%	2	1	8	1	5	4	5	6	9	7	8

+ Data collection for 2018 and 2019 is only partially complete. Italicized estimates may change in the future if more reports of deaths are received.


Source: U.S. Consumer Product Safety Commission / EPHA.

CPSC Death Certificate File, CPSC In-Depth Investigation File, CPSC Injury or Potential Injury Incident File,  
National Center for Health Statistics Mortality File, 2009–2019

Note: Reported annual estimates and estimated averages and percentages may not add to subtotals or totals due to rounding.

Figure 2 graphically illustrates the relationship between the time of year and the estimated number of CO poisoning deaths from 2009 through 2019. The total estimated number of CO poisoning deaths is presented on the radar graph by month of death. The shaded area represents the estimated total number of deaths for the 11-year period, distributed by each month of a year. Notably, more CO deaths occur in the cold months, particularly November, December, January, and February, than in warm months. Fatalities increase as the winter months continue, until a slight drop off in February before the spring months come. Conversely, as time gets deeper into the warmer months, the number of deaths decreases, with the lowest number of fatalities occurring in July and August.

**Figure 2: Estimated Number of Consumer Product-Related Carbon Monoxide Deaths by Month of Death, 2009–2019**



Source: U.S. Consumer Product Safety Commission/EPHA. CPSC Death Certificate File, CPSC In-Depth Investigation File, CPSC Injury or Potential Injury Incident File, National Center for Health Statistics Mortality File, 2009–2019.

# Victim Demographics from Non-Fire Carbon Monoxide Poisoning Deaths Associated with the Use of Consumer Products

## Age of Victim

Table 7 shows the estimated number of CO poisoning deaths categorized by victim age for the 11 most recent years of data (2009–2019). From the data, it appears that consumer product-related CO deaths are skewed toward older individuals. For the 3 most recent years (2017–2019), children younger than 15 years of age accounted for an annual average of 4 percent (an estimated 9 deaths out of 216) of the yearly CO poisoning deaths, while this age group represents an average of about 19 percent of the U.S. population. For the same time frame, deaths among adults 45 years and older accounted for more than 66 percent (143 of 216), while this age group represented about 42 percent of the U.S. population. Statistical tests confirm (see Appendix C for p-values) the significance in the age-related differences in CO poisoning deaths. Percentages of children below the age of 15, as well as individuals 15 to 24 years old were each identified as statistically significantly below population estimates. Conversely, percentage of CO deaths among individuals 45 to 65, and those over age 65, were identified as statistically greater than their population representation.

**Table 7: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Age of Victim, 2009–2019**

Age	2017–2019 <sup>+</sup>		Estimated Percentage of U.S. Population <sup>@</sup>	Annual Estimate										
	Average Estimate	Average Percent		2009	2010	2011	2012	2013	2014	2015	2016	2017	2018 <sup>+</sup>	2019 <sup>+</sup>
<b>Total</b>	<b>216</b>	<b>100%</b>	<b>100%</b>	<b>148</b>	<b>159</b>	<b>163</b>	<b>137</b>	<b>146</b>	<b>164</b>	<b>172</b>	<b>178</b>	<b>188</b>	<b>210</b>	<b>250</b>
Under 5	3	2%	6%	3	1	*	1	*	2	*	1	2	4	4
5 - 14	6	3%	13%	2	1	4	4	5	7	17	6	9	1	7
15 - 24	13	6%	13%	14	12	9	6	11	8	15	6	17	7	14
25 - 44	51	24%	27%	43	39	36	37	34	35	45	54	55	55	43
45 - 64	89	41%	26%	59	69	63	56	62	67	65	83	55	89	124
65 and over	54	25%	16%	27	36	52	32	36	44	31	29	51	54	58

+ Data collection for 2018 and 2019 is only partially complete. Italicized estimates may change in the future if more reports of deaths are received.

\* No reports received by CPSC staff.

@ Based on estimated U.S. population statistics for the 3- year average (2017-2019). U.S. Census Bureau.

Source: U.S. Consumer Product Safety Commission/EPHA.

CPSC Death Certificate File, CPSC In-Depth Investigation File, CPSC Injury or Potential Injury Incident File, National Center for Health Statistics Mortality File, 2009-2019.

U.S. Census Bureau, Population Division. Annual Estimates of the Resident Population by Sex, Age, Race, and Hispanic Origin for the United States and States: April 1, 2010 to July 1, 2019. June 2020.

Note: Reported annual estimates and estimated averages and percentages may not add to subtotals or totals due to rounding.

## Gender of Victim

Table 8 presents the distribution of estimated CO deaths categorized by gender. In 2019, 77 percent of CO poisoning victims were males, and 23 percent were females. These percentages varied slightly from year to year over the 11 years of this report. However, every year there were many more male CO deaths than female. For 2017 through 2019, the average percentage of male CO victims was 76 percent, and the average percentage of female victims was 24 percent. By contrast, about 49 percent of the U.S. population is male, and 51 percent of the U.S. population is female.<sup>3</sup> The gender-related differences in CO poisoning deaths were confirmed to be statistically significant (p-value < 0.0001).

**Table 8: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Gender of Victim, 2009-2019**

Gender	2017–2019 <sup>+</sup>		Estimated Percentage of U.S. Population*	Annual Estimate										
	Average Estimate	Average Percent		2009	2010	2011	2012	2013	2014	2015	2016	2017	2018 <sup>+</sup>	2019 <sup>+</sup>
<b>Total</b>	<b>216</b>	<b>100%</b>	<b>100%</b>	<b>148</b>	<b>159</b>	<b>163</b>	<b>137</b>	<b>146</b>	<b>164</b>	<b>172</b>	<b>178</b>	<b>188</b>	<b>210</b>	<b>250</b>
Male	<b>165</b>	<b>76%</b>	49%	109	121	111	92	124	127	125	140	138	<i>164</i>	<i>192</i>
Female	<b>51</b>	<b>24%</b>	51%	39	38	52	45	22	37	48	38	50	<i>46</i>	<i>57</i>

+ Data collection for 2018 and 2019 is only partially complete. Italicized estimates may change in the future if more reports of deaths are received.

\* Based on estimated U.S. population statistics for the 3-year average (2017-2019).

Source: U.S. Consumer Product Safety Commission/EPHA.

CPSC Death Certificate File, CPSC In-Depth Investigation File, CPSC Injury or Potential Injury Incident File, National Center for Health Statistics

Mortality File, 2009–2019.

U.S. Census Bureau, Population Division. Annual Estimates of the Resident Population by Sex, Age, Race, and Hispanic Origin for the United States and States:

April 1, 2010 to July 1, 2019. June 2020

Note: Reported annual estimates and estimated averages and percentages may not add to subtotals or totals due to rounding.

<sup>3</sup> Three-year average, 2017 to 2019, from June 2020 U.S. Census estimates of the U.S. population.

## Victim Race/Ethnicity

Table 9 provides a summary of CO fatality victims characterized by race/ethnicity for the years 2009 through 2019. Because of the growing proportion of people of Hispanic descent, Hispanic victims were categorized separately, irrespective of their race. Estimates of the percentage of the U.S. population categorized into the various race/ethnicity groupings were based on single-race characterizations, as represented in the U.S. Census Bureau reports. Non-Hispanic individuals reported as multiracial are included in the *Unknown/Other/Mixed* category.

The estimated percentages of the 2017 through 2019 annual average CO deaths demonstrated some race/ethnicity-based differences in CO poisoning deaths that were statistically significant ( $p$ -value = 0.0400). When looked at as one race/ethnicity versus the rest, there was a statistically significant difference between the number of White victims of CO poisoning (approximately 69 percent of all CO poisoning deaths) and the resident White population (about 60 percent of the U.S. population), the  $p$ -value of this comparison was 0.0489. CO fatalities among Black and African American represented 11 percent of all CO fatalities, while their representation in the U.S. population is 13 percent. The difference is not determined to be statistically significant. However, in prior years of this report, the proportion of Black or African American victims has been determined to be statistically significantly greater than the resident population. It is unclear whether this is an anomaly in the recent data or an actual change.

Additionally, as has been seen before, the proportion of the CO poisoning fatality victims who were of Hispanic ethnicity (approximately 11%) was below the percentage of Hispanics in the U.S. population (about 18%), where the  $p$ -value was 0.0094. Among other races/ethnicities, no statistically significant differences were observed.

**Table 9: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Race/Ethnicity, 2009–2019**

Race/ Ethnicity	2017–2019 <sup>+</sup>		Estimated Percentage of U.S. Population <sup>@</sup>	Annual Estimate										
	Average Estimate	Average Percent		2009	2010	2011	2012	2013	2014	2015	2016	2017	2018 <sup>+</sup>	2019 <sup>+</sup>
<b>Total</b>	<i>216</i>	<i>100%</i>	100%	148	159	163	137	146	164	172	178	188	<i>210</i>	<i>250</i>
White <sup>4</sup>	<i>149</i>	<i>69%</i>	60%	93	82	106	82	86	108	109	118	121	<i>153</i>	<i>173</i>
Black or African American <sup>4</sup>	23	11%	13%	20	43	38	31	35	26	47	32	24	18	26
Hispanic (All races)	25	11%	18%	11	18	9	11	13	18	14	13	19	26	29
Asian/Pacific <sup>1</sup>	5	2%	6%	3	4	3	5	7	6	*	7	7	4	5
American Indian <sup>2</sup>	5	2%	1%	1	5	1	*	1	1	*	1	7	4	5
Unknown/Other /Mixed <sup>3</sup>	9	4%	2%	19	8	6	7	5	5	3	6	9	6	11

+ Data collection for 2018 and 2019 is only partially complete. Italicized estimates may change in the future if more reports of deaths are received.

\* No reports received by CPSC staff.

@ Based on estimated U.S. population statistics for the 3- year average (2017-2019). U.S. Census Bureau.

1 Includes Asian, Pacific Islander, and Native Hawaiian

2 Includes American Indian, Native American, and Native Alaskan

3 Includes non-Hispanic Unknown races, Other races, and Multiple races

4 Only includes non-Hispanic ethnicities.

Source: U.S. Consumer Product Safety Commission/EPHA.

CPSC Death Certificate File, CPSC In-Depth Investigation File, CPSC Injury or Potential Injury Incident File,  
National Center for Health Statistics Mortality File, 2009-2019.

U.S. Census Bureau, Population Division. Annual Estimates of the Resident Population by Sex, Age, Race, and Hispanic Origin for the United States and  
States: April 1, 2010 to July 1, 2019. June 2020.

Note: Reported annual estimates and estimated averages and percentages may not add to subtotals or totals due to rounding.

## Population Density of Place of Death

Table 10 provides a breakdown of the CO poisoning deaths characterized by population density of the incident location. The table is presented as three sections: (1) incidents occurring at all locations; (2) incidents occurring in locations identified as a permanent home (e.g., house, apartment, mobile home); and (3) incidents occurring only in non-home locations (e.g., camper trailer, tent, motel room). Please note that “Home Locations” and “Non-Home Locations” sum to “All Locations.”

All fatal incidents were designated as occurring in one of four rural/urban categories, based on the Rural-Urban Commuting Area (RUCA) codes developed by the Economic Research Service (ERS) of the U.S. Department of Agriculture (USDA) in conjunction with the Center for Rural Health, School of Medicine and Health Sciences, University of North Dakota. The categories are based on theoretical concepts used by the U.S. Office of Management and Budget (OMB) to define county-level metropolitan and micropolitan areas.<sup>4</sup> This 21-category classification system is based on measures of population density, urbanization, and daily commuting. The OMB methodology is based on a county-level delineation. ERS refined the methodology by applying it to smaller census tracts. ERS further delineated the characterization by cross-referencing each zip code in the United States to its RUCA code classification.<sup>5</sup> The update of the RUCAs to version 3.1 was developed by Center for Rural Health, School of Medicine and Health Sciences, University of North Dakota and ERS and is funded by the U.S. Department of Health and Human Services, Health Resources and Services Administration, Office of Rural Health Policy, and the USDA Economic Research Service. The zip code cross-reference was used to characterize each of the CO deaths into one of four broad categories: Urban Core, Sub-Urban, Large Rural Town, and Small Town/Rural Isolated. The RUCA codes are updated approximately once every 10 years. The most recent update applicable to years addressed in this report was for the year 2010. It is unlikely that there would be a substantial change in the urban-rural population distribution between 2010 and the more recent 3-year period average of 2017 through 2019.

Table 10 also includes the estimated percentage of the U.S. population, per population density designation category. As can be seen in the *All Locations* section, the estimated average percentage of CO deaths during the 3-year period 2017 through 2019, in urban locations (50%), is smaller than the percentage of the U.S. population living in urban core locations (73%). The difference is offset by the larger percentages in the other three categories: sub-urban locations (24% versus 15% of the U.S. population), large rural town locations (9% versus 6%), and small town/rural isolated locations (15% versus 5%). CO deaths that occurred in small town/rural isolated locations were nearly three times the percentage of the U.S. population living in these isolated locations. Additionally, due to lack of detail in some of the death certificates that CPSC receives, the exact location of some incidents (6%) could not be ascertained. The 2017 through 2019 data do not show a distinct difference between Home Locations and Non-Home Locations.

---

<sup>4</sup> OMB BULLETIN NO. 13-01: Revised Delineations of Metropolitan Statistical Areas, Micropolitan Statistical Areas, and Combined Statistical Areas, and Guidance on Uses of the Delineations of these Areas. February 28, 2013.

<sup>5</sup> Version 3.10 of the ZIP code Rural-Urban Commuting Areas (RUCAs) geographic taxonomy, August 4, 2014. Center for Rural Health, University of North Dakota School of Medicine and Health Sciences. Comparable data presently available from [USDA ERS - Rural-Urban Commuting Area Codes](#).

**Table 10: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Population Density of Place of Death, 2009–2019**

RUCA Population Density Designation	2017–2019+		Estimated Percentage of US Population@	Annual Estimates										
	Average Estimate	Average Percent		2009	2010	2011	2012	2013	2014	2015	2016	2017	2018+	2019+
<b>All Locations</b>	216	100%	100%	148	159	163	137	146	164	172	178	188	210	250
Urban Core	108	50%	73%	78	94	95	79	84	73	77	108	103	107	114
Sub-Urban	52	24%	15%	42	33	33	25	27	34	41	25	39	51	65
Large Rural Town	19	9%	6%	10	25	14	9	12	19	14	20	13	24	21
Small Town/Rural Isolated	32	15%	5%	18	7	18	19	23	32	39	22	33	27	37
Unknown Location	4	2%	-	*	*	2	6	1	6	1	2	*	*	13
<b>Home Locations</b>	162	75%	100%	117	135	137	113	121	127	131	150	150	148	188
Urban Core	83	38%	73%	66	88	78	71	73	63	62	92	83	76	89
Sub-Urban	39	18%	15%	30	24	28	20	24	29	39	21	33	32	51
Large Rural Town	15	7%	6%	10	19	14	6	7	14	11	18	9	18	17
Small Town/Rural Isolated	25	12%	5%	11	4	15	11	15	21	18	18	27	22	27
Unknown Location	1	1%	-	*	*	2	5	1	*	*	*	*	*	4
<b>Non-Home Locations</b>	53	25%	100%	30	24	26	24	26	37	41	28	38	62	62
Urban Core	26	12%	73%	11	6	18	7	11	11	14	16	21	32	26
Sub-Urban	13	6%	15%	12	8	5	5	2	5	1	3	6	20	14
Large Rural Town	5	2%	6%	*	6	*	2	5	5	3	2	5	6	4
Small Town/Rural Isolated	7	3%	5%	7	4	3	7	8	11	22	4	6	5	9
Unknown Location	3	1%	-	*	*	*	1	*	6	1	2	*	*	9

+ Data collection for 2018 and 2019 is only partially complete\* No reports received by CPSC staff.

@ Estimated 2010 U.S. population categorized by Rural Urban Commuting Area (RUCA 3.1) designation. U.S. population estimates by RUCA classification were determined by cross-referencing the Center for Rural Health, School of Medicine and Health Sciences, University of North Dakota/Economic Research Service, Department of Agriculture RUCA3.1 zip code table with the 2010 U.S. Census population estimates by zip code area.019 is only partially complete. Italicized estimates may change in the future if more reports of deaths are received.

Source: U.S. Consumer Product Safety Commission / EPA.

CPSC Death Certificate File, CPSC In-Depth Investigation File, CPSC Injury or Potential Injury Incident File, National Center for Health Statistics Mortality File, 2009– 2019

Center for Rural Health, University of North Dakota School of Medicine and Health Sciences, ZIP code RUCA Version 3.10

Note: Reported annual estimates and estimated averages and percentages may not add to subtotals or totals due to rounding.

## Geographical Region of Incident

Table 11 provides a breakout of the CO poisoning deaths characterized by geographic region where the incident occurred. As the table reflects, for the 3 most recent years (2017 to 2019), CO deaths in some of the regions appear to be different from what would be expected based on the percentage of the U.S. population living in these regions. This may indicate that geographic location influences the likelihood of fatal CO poisoning incidents; however, these results may be influenced due to incompleteness of the estimates for a few states. The regional estimates and proportions for recent years, therefore, are not assessed for statistical significance.

The states that comprise each of the regions are set forth in Appendix D.

**Table 11: Estimated Non-Fire Carbon Monoxide Poisoning Deaths by Geographical Region of Incident, 2009–2019**

Region <sup>‡</sup>	2017–2019 <sup>+</sup>		Estimated Percentage of US Population <sup>@</sup>	Annual Estimates										
	Average Estimate	Average Percent		2009	2010	2011	2012	2013	2014	2015	2016	2017	2018+	2019 <sup>+</sup>
<b>Total</b>	<b>216</b>	<b>100%</b>	<b>100%</b>	<b>148</b>	<b>159</b>	<b>163</b>	<b>137</b>	<b>146</b>	<b>164</b>	<b>172</b>	<b>178</b>	<b>188</b>	<b>210</b>	<b>250</b>
<b>Northeast</b>	<b>27</b>	<b>13%</b>	<b>17%</b>	<b>14</b>	<b>23</b>	<b>43</b>	<b>25</b>	<b>34</b>	<b>37</b>	<b>30</b>	<b>21</b>	<b>11</b>	<b>30</b>	<b>41</b>
New England	9	4%	5%	5	5	16	1	14	8	13	8	4	12	10
Middle Atlantic	19	9%	13%	9	18	27	24	20	29	17	13	7	18	31
<b>South</b>	<b>70</b>	<b>32%</b>	<b>38%</b>	<b>55</b>	<b>55</b>	<b>55</b>	<b>55</b>	<b>43</b>	<b>42</b>	<b>62</b>	<b>55</b>	<b>67</b>	<b>74</b>	<b>69</b>
East South Central	14	6%	6%	19	12	13	7	3	9	11	10	10	20	13
South Atlantic	39	18%	20%	13	26	23	31	20	21	29	27	36	43	38
West South Central	17	8%	12%	23	17	19	17	20	12	22	18	21	11	18
<b>Midwest</b>	<b>58</b>	<b>27%</b>	<b>21%</b>	<b>48</b>	<b>49</b>	<b>33</b>	<b>31</b>	<b>48</b>	<b>40</b>	<b>44</b>	<b>69</b>	<b>61</b>	<b>53</b>	<b>59</b>
East North Central	36	17%	14%	28	40	27	26	27	22	27	44	40	26	43
West North Central	21	10%	7%	20	10	6	5	21	18	17	25	21	27	16
<b>West</b>	<b>62</b>	<b>29%</b>	<b>24%</b>	<b>31</b>	<b>31</b>	<b>32</b>	<b>25</b>	<b>22</b>	<b>44</b>	<b>37</b>	<b>33</b>	<b>50</b>	<b>54</b>	<b>81</b>
Mountain	27	13%	7%	16	11	9	13	8	26	14	18	12	36	33
Pacific	35	16%	16%	14	20	23	12	14	18	23	15	38	18	48

‡ Region designation is based on U.S. Census Bureau reporting practices. See Appendix D for identification of specific regional designation of state of occurrence.

+ Data collection for 2018 and 2019 may be only partially complete due to apparently incomplete reporting from some states. Italicized estimates may change in the future if more reports of deaths are received.

@ Based on estimated U.S. population statistics for the 3-year average (2017-2019).

Source: U.S. Consumer Product Safety Commission/EPHA.

## Appendix A: Methodology

This appendix describes the data sources and methodology used to compute the national estimate of non-fire carbon monoxide (CO) poisoning deaths associated with the use of consumer products and the estimates by product, victim age, and incident location.

All death certificates filed in the United States are compiled by the National Center for Health Statistics (NCHS) into a multiple cause-of-mortality data file. The NCHS Mortality File contains demographic and geographic information, as well as the International Statistical Classification of Diseases and Related Health Problems codes for the underlying cause of death. Data are compiled in accordance with the World Health Organization instructions, which request that member nations classify causes of death by the current Manual of the International Statistical Classification of Diseases and Related Health Problems. The International Classification of Diseases, Tenth Revision (ICD-10) was implemented in 1999. Although the NCHS data contain cause-of-death codes that are helpful in identifying deaths due to CO poisoning, the records do not contain any narrative information that might indicate the involvement of a consumer product.

CPSC staff purchases death certificates from the 50 states, New York City, the District of Columbia, and some U.S. territories. Specifically, CPSC staff purchases death certificates with certain cause-of-death codes for which a high probability exists that consumer products are involved. In addition to the cause-of-death codes and demographic and geographic information, the death certificate contains information about the incident location and a brief narrative describing the incident. Any references to consumer products are usually found in these narratives. As resources allow, CPSC staff conducts follow-up In-Depth Investigations (IDIs) on selected deaths to confirm and expand upon the involvement of consumer products. These data from CPSC complement the NCHS mortality data.

ICD-10 classifies deaths associated with CO poisoning with the codes listed below. The focus of this report is accidental CO poisoning deaths, and the report concentrates on deaths coded as X47 and Y17. Deaths coded under Code X67, intentional CO poisonings, are excluded from this analysis.

## ICD-10 Code    Definitions

X47	<b>Accidental</b> – Poisoning by and exposure to other gases and vapors.  Includes: carbon monoxide, lacrimogenic gas, motor (vehicle) exhaust gas, nitrogen oxides, sulfur dioxide, utility gas.
X67	<b>Intentional</b> – Poisoning by and exposure to other gases and vapors.  Includes: carbon monoxide, lacrimogenic gas, motor (vehicle) exhaust gas, nitrogen oxides, sulfur dioxide, utility gas.
Y17	<b>Undetermined intent</b> – Poisoning by and exposure to other gases and vapors.  Includes: carbon monoxide, lacrimogenic gas, motor (vehicle) exhaust gas, nitrogen oxides, sulfur dioxide, utility gas.

The first step in compiling the annual estimates is computing the total estimates of CO poisoning deaths associated with consumer products. The CPSC's Death Certificate (DTHS) File and the CPSC's Abbreviated Death Certificate (ABDT) File were searched for cases associated with ICD-10 codes X47 and Y17.

Each case in the CPSC's DTHS File that was coded as X47 or Y17 was reviewed by an analyst and categorized as in-scope, out-of-scope, or source of CO unknown or questionable. In-scope cases are unintentional, non-fire CO poisoning deaths associated with a consumer product under the jurisdiction of the CPSC. Out-of-scope cases are cases that involve CO sources that are not under the jurisdiction of the CPSC, fire- or smoke-related exposures, or intentional CO poisonings. Examples of out-of-scope cases include poisonings due to gases other than CO (*i.e.*, natural gas, ammonia, butane); motor vehicle exhaust- or boat exhaust-related poisonings; and work-related exposures. The source of CO was classified as unknown or questionable in cases where a consumer product was possibly associated with the incident, but the exact source of CO was unknown.

The CPSC's ABDT File contains death certificates for CO poisonings (X47 and Y17) that involve motor vehicle exhaust, cases where the source of the CO is unknown, or where the death certificate does not mention a consumer product. Other examples of cases that may appear in the abbreviated file are cases associated with farm accidents, smoke inhalation from a structural fire, or other gas poisonings. Occasionally, newer information from CPSC IDIs may be matched with ABDT cases that were originally classified as having no known source or did not mention a consumer product. If information from IDIs indicated that an ABDT case should be considered in scope, then it was included with the DTHS database files. For 2008, 2009, 2010, and 2011, no ABDT records were reclassified as in scope. From 2012 through 2017, nine cases were reclassified: three cases for 2012; one case for 2013; four cases for 2014; two cases in 2015; one case in 2016; two cases in 2017. No cases were reclassified in 2018. In 2019, only one ABDT record was reclassified as in scope.

In 2016, and to a slightly lesser extent in 2017, the way the state of Texas designated death certificates with the Y17 code seems to have changed. Before 2016, the maximum number of

Y17-coded death certificates from any individual state was 21 (coincidentally, Texas in 2013). In 2016, CPSC received 56 Y17-coded death certificates from Texas, and 129 from the entire country. In 2016, Michigan, the second highest number of Y17, had 13. In 2017, death certificates from Texas with the Y17 code dropped to 34 but were still much higher than any state for any year. In 2017, the second highest number of Y17-coded death certificates was only six by Oklahoma and Oregon. NCHS records indicate 94 Y17s in 2016, and 85 in 2017. For these two years, CPSC has 90 Y17-coded death certificates from Texas, more than the rest of the country combined. Clearly, in 2016 and 2017, some discrepancy exists with the way Texas codes Y17 death certificates compared to the rest of the states in the country. As noted in the prior year's report, it appeared as though there were many 2018 death certificates missing from the CPSC database as of the search date for that year's report. That report anticipated additional collection (from particular states), but the expected missing reports have gone unreported without any change from July 2021 through September 2022. CPSC data for 2019 also appears to be incomplete from all states, even if most are complete. With reporting for 2018 and 2019 still incomplete, no comparable adjustments are made for 2019, or changes to the prior calculations for previous years, in the absence of the anticipated additional death certificate reporting from some states for those prior years.

Thus far, Texas, Washington, and Wisconsin have reported considerably fewer relevant deaths for the years 2018 and 2019 than typical of previous years. Therefore, the death certificate data available to CPSC for 2018 and 2019, likely underrepresents the distribution of deaths among those three states (*i.e.*, Texas, Washington, and Wisconsin). Estimation methodologies in this report generally assume randomized non-reporting, which does not appear to represent an accurate assumption of distribution at the individual state level for 2018 and 2019. To a lesser degree, incomplete estimates *by region* may also merit some skepticism for the incomplete years 2018 and 2019. Despite these caveats, the incomplete estimates for 2018 and 2019 nevertheless, may provide valuable information, in assessing the U.S. as a whole.

To compensate for the apparent anomalies in 2016 and 2017 only, this report maintains the most recent report's substitution of the average yearly number of Y17 reports from the prior 10 years for Texas, in place of the 2016 and 2017 count of Texas Y17s in the scaling calculations. The average number of Y17-coded death certificates from the previous 10 years is 7.6. However, no similar compensation appears appropriate yet for 2018 and 2019.

**Table A.1: Initial Categorization for 2019 Data**

ICD-10 Code	NCHS Total	CPSC DTHS File & ABDT File				Number of Cases to be Imputed
		In-Scope	Unknown Scope	Out-of-Scope	Total	
X47	994	189	51	572	812	233
Y17	112	3	9	108	120	1
<b>Total</b>	<b>1,106</b>	<b>192</b>	<b>60</b>	<b>680</b>	<b>932</b>	<b>234</b>

<sup>1</sup> "NCHS Total" cases, minus "Total in CPSC Database," plus "Unknown Scope" from DTHS.

Source: U.S. Consumer Product Safety Commission/EPHA.

CPSC Death Certificate File, CPSC In-Depth Investigation File, Abbreviated Death Certificate File, National Center for Health Statistics Mortality File, 2019.

The proportion of death certificates found in the CPSC database associated with non-fire, unintentional X47 or Y17 deaths and associated with consumer products was applied to the NCHS totals to calculate the total estimated number of non-fire CO poisoning deaths associated with consumer products. In theory, the NCHS totals comprise all death certificates in the United States, and the same proportion of in-scope cases should exist in the death certificates that are missing from the combined CPSC Death Certificate and Abbreviated Death Certificate files or are from an unknown source. Applying the proportion of in-scope cases to the NCHS database totals, therefore, should provide an estimate of in-scope cases nationwide. This was done in the following way for ICD-10 codes X47 and Y17, separately:

1. The number of in-scope deaths in the CPSC's two death certificate files coded under the specific ICD10 code that were associated with an accidental non-fire CO poisoning and a consumer product were identified ( $n_1$ ).
2. The total number of deaths in the CPSC's Death Certificate File and the Abbreviated Death Certificate File coded under the specific ICD10 code were summed separately, excluding cases with an unknown or highly questionable source ( $n_2$ ).
3. The total number of deaths in the NCHS data associated coded under the specific ICD10 code was counted ( $n_3$ ).
4. The estimate of the number of non-fire CO poisoning deaths associated with consumer products under the specific ICD-10 code was calculated, using the formula:

$$N = (n_1 / n_2) * n_3$$

The proportion ( $n_1/n_2$ ) represents the number of in-scope cases found in the CPSC's files, divided by the total of in-scope and out-of-scope cases.

5. The estimates of the number of non-fire CO poisoning deaths associated with consumer products under the specific ICD10 codes were summed to calculate the total estimate of non-fire CO poisoning deaths.

$$\text{Total Estimate} = N_{X47} + N_{Y17}$$

The ratio ( $n_3 / n_2$ ) represents the weighting factor used to calculate the annual estimates. The CPSC's Death Certificate File does not contain death certificates for all deaths listed in the NCHS file; therefore, a weighting factor was calculated to account for death certificates that are missing. The weighting factor allows the computation of national estimates of CO deaths by consumer products and by other characteristics collected by CPSC about each death.

Table A.2 contains the values for the variables used in the calculation, as well as the final computed 2019 estimates of CO poisoning deaths.

**Table A.2: Calculation Detail of the Final Computed 2019 Estimate of Non-Fire CO Poisoning Deaths Associated with Consumer Products**

Variable	ICD-10 Code	
	X47	Y17
$n_1$	189	3
$n_2$	$812 - 51 = 761$	$120 - 9 = 111$
$n_3$	994	112
<i>Weighting Factor (<math>n_3 / n_2</math>)</i>	1.3062	1.0090
N	246.8673	3.027
Total Estimate	$\{246.8673 + 3.027 = 249.8943 \sim 250\}$	

Source: U.S. Consumer Product Safety Commission/EPHA. CPSC Death Certificate File, CPSC In-Depth Investigation File, Abbreviated Death Certificate File, National Center for Health Statistics Mortality File 2019.

Death certificates received by NCHS are routinely checked for accuracy of state personnel-identified ICD-10 coding. On occasion, NCHS staff will correct codes before entering the data into their databases. CPSC staff has no way of correcting CPSC records to mesh with NCHS records. CPSC receives death certificate facsimiles or electronic death certificates directly from the states before any possible corrections are deemed necessary per NCHS procedures. Consequently, there may be slight discrepancies between final NCHS counts and CPSC records. For this report, CPSC staff has assumed that, over time, the number of death certificates with ICD-10 codes changed by NCHS staff to the codes of interest (X47 and Y17), would equal approximately those changed to codes other than X47 or Y17, thereby having little long-term effect on the estimates.

Table A.3 shows the weighting factors used to calculate the estimates for the years 2009–2019, based on the information available to CPSC staff.

**Table A.3: CO Fatality Cases and Weighting Factors Used to Calculate the Estimates for the Years 2009–2019**

Year	NCHS Total	Total in CPSC Databases*	In-Scope Cases	Weighting Factor
2009				
X47	734	769	145	1.0000
Y17	72	52	2	1.3846
2010				
X47	675	567	125	1.1905
Y17	98	68	7	1.4412
2011				
X47	786	730	143	1.0767
Y17	89	76	8	1.1711
2012				
X47	736	591	109	1.2453
Y17	114	84	1	1.3571
2013				
X47	704	608	123	1.1579
Y17	76	60	3	1.2667
2014				
X47	803	679	137	1.1826
Y17	106	61	1	1.7377
2015				
X47	847	665	134	1.2737
Y17	91	53	1	1.7170
2016				
X47	921	822	154	1.1204
Y17	94	72.6	4	1.2948
2017				
X47	936	770	150	1.2156
Y17	85	75.6	5	1.1243
2018				
X47	896	730	164	1.2274
Y17	106	97	8	1.0928
2019				
X47	994	761	189	1.3062
Y17	112	111	3	1.0090

\* This is the total number of deaths in the Death Certificate File and Abbreviated Death Certificate File, excluding deaths associated with an unknown or questionable source of CO.

Source: U.S. Consumer Product Safety Commission/EPHA. CPSC Death Certificate File, CPSC In-Depth Investigation File, Abbreviated Death Certificate File, National Center for Health Statistics Mortality File, 2009–2019.

Incidents with unknown or highly questionable CO sources were excluded from the denominator (the number of deaths in the CPSC databases) of the weighting factor. The group of cases with unknown or highly questionable sources was assumed to contain the same proportion of cases associated with a consumer product as the group of cases within the CPSC database with known CO sources (this is the same assumption that is made for cases where the death certificate is missing). To include these cases within the denominator assumes that these cases can be classified as in-scope or out-of-scope cases when their scope status is unknown.

Therefore, for weighting purposes, cases with unknown or questionable sources were treated in the same way as missing cases.

In-scope cases were examined further to determine which product was associated with the incident. Additional information on the CO deaths was obtained from review of the CPSC's IDI File.


Reports of non-fire CO poisoning deaths were retrieved from the DTHS and ABDT files, based on the following criteria: date of death between 1/1/2009 and 12/31/2019, and ICD-10 code of X47 or Y17. Death certificates entered in the CPSC's database before September 1, 2022, were included in this analysis. Whenever possible, each CO death was reviewed and coded by the author, according to the consumer product and type of fuel involved, incident location, and whether multiple deaths were associated with the same incident.

In Table 1 of this report, the *Heating Systems* category includes CO poisoning deaths from subcategories for furnaces and boilers (combined under the heading of *Furnaces*), vented floor and wall heaters, unvented room/space heaters, unvented portable heaters, and other miscellaneous heating systems. Each subcategory is further delineated by fuel type used. Deaths associated with charcoal burned alone and in the absence of an appliance (e.g., in a pail or in the sink) were presented with *Charcoal/Charcoal Grills*, even though this practice typically is done for heating purposes. Examples of products historically included in the *Other Products* category include LP gas refrigerators and gas pool heaters. LP gas grill, LP fish cooker, and other LP gas portable cooking appliance incidents are classified in the *Grills, Camp Stoves* category. Deaths where multiple fuel-burning products were used simultaneously, such that a single source of the fatal CO could not be determined, were classified under *Multiple Products*. *Engine-Driven Tools* included generators and power gardening equipment, such as power lawn mowers, garden tractors, concrete cutters, gasoline-powered water pumps, and snow blowers. Generators that were original equipment installed on a recreational vehicle (RV), trailer, camper, or boat were considered out of scope because they are likely outside the jurisdiction of the CPSC.

# Appendix B: National Estimates and Mortality Rates of Consumer Product-Related CO Poisoning Deaths, 1980 to 2018

Figure B.1 below graphically suggests a trend of the estimated CO deaths from 1980 to 2019. Before the implementation of the ICD-10 coding in 1999, the estimated number of non-fire, consumer product-related CO poisoning deaths decreased from the early 1980s to the late 1990s, from a high of 340 in 1982, to a low of 180 in both 1997 and 1998. In 1999, there were an estimated 108 consumer product-related CO deaths, well below the estimated 180 deaths in each of the two previous years. The difference may be due, in part, to the change from ICD-9 coding to ICD-10 coding, where product identification could be assessed more accurately. As can be seen in the graph below, 2018 was the first year since ICD-10 was implemented in 1999 to exceed 200 CO fatalities before increasing to an estimated 250 such fatalities in 2019. However, some part of this increase is due to population growth. According to the U.S. Census, the U.S. population grew by more than 17 percent between 1999 and 2019.

**Figure B.1: Estimated Non-Fire CO Poisoning Deaths Associated with Consumer Products: 1980–2019**



**Implementation of ICD-10.**

Source: U.S. Consumer Product Safety Commission/EPHA. CPSC Death Certificate File, CPSC Injury or Potential Injury Incident File, CPSC In-Depth Investigation File, 2009–2019.

## Estimated 3-Year CO Mortality Trends

Table B.1 presents the annual estimates from 1980 to 2019, and the 3-year average mortality rates associated with each year, where 3 years of data were available. The 3-year average mortality rate is presented in the table for the mid-point year. The estimated 3-year average mortality rate decreased from the 1982 high of 14.02 per 10 million population, to a 3-year average rate of 4.34 per 10 million in 2000, a reduction of 69 percent. Subsequently, the 3-year average rate increased annually through 2006, to a rate of 6.21. Since 2006, the rate slowly dropped to the 2013 estimate of 4.71, before reversing the trend and rising in the 2018 estimate to a rate of 6.61. This 2018 rate estimate exceeds the 2006 estimate of 6.21, which still included the effects of the devastation of Hurricane Katrina and other 2005 hurricanes.

**Table B.1: Estimated Non-Fire Carbon Monoxide Poisoning Death Rates Associated with Consumer Products, 1980–2019**

Year	Estimate	U.S. Population Estimates (thousands)	3-Year Average Mortality Rate per 10 Million Population
1980	282	227,225	
1981	311	229,466	13.55
1982	340	231,664	14.02
1983	323	233,792	13.38
1984	275	235,825	12.47
1985	284	237,924	11.19
1986	240	240,133	10.49
1987	232	242,289	9.77
1988	238	244,499	10.44
1989	296	246,819	10.49
1990	243	249,623	10.53
1991	250	252,981	9.27
1992	211	256,514	8.77
1993	214	259,919	8.31
1994	223	263,126	8.08
1995	201	266,278	8.02
1996	217	269,394	7.40
1997	180	272,647	7.05
1998	180	275,854	5.66
1999*	108	279,040	5.09
2000	138	282,172	4.34
2001	121	285,082	5.15
2002	181	287,804	5.27
2003	153	290,326	5.76

2004	168	293,046	5.81
2005	190	295,753	6.06
2006	180	298,593	6.21
2007	186	301,580	6.01
2008	178	304,375	5.61
2009	148	307,007	5.27
2010	159	309,338	5.06
2011	163	311,644	4.91
2012	137	313,993	4.74
2013	146	316,235	4.71
2014	164	318,857	5.05
2015	172	321,419	5.34
2016	178	323,128	5.55
2017	188	325,719	5.90
2018	210	326,838	6.61
2019	250	328,240	

Note: The 3-year average mortality rate is reported at the mid-point year.

\* The Tenth Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10) was implemented.

Source: U.S. Consumer Product Safety Commission/EPHA.

U.S. Census Bureau, Population Division. Annual Estimates of the Resident Population by Sex, Age, Race, and Hispanic Origin for the United States and States: April 1, 2010 to July 1, 2019. June 2020.

Before implementation of ICD-10 in 1999, generating estimates for an important category of products—generators and other engine-driven tools—was not possible.<sup>6</sup> With the advent of ICD-10 coding, generating estimates of deaths associated with generators and other engine-driven tools is now possible. Table B.2 presents a summary of the mortality rates associated with generators, which steadily increased from 1999 through 2006, but retracted until 2011, from the previous 2006 high point. However, the rate generally increased after 2013, with the most recent 3-year average for 2016 to 2018, as the highest level (2.85) so far, exceeding the previously highest 2.69 rate in 2006, which included the Hurricane Katrina impact of 2005. This most recent 3-year average mortality rate range for generators alone is more than five times greater than the 3-year average rate in 2000.

---

<sup>6</sup> See Appendix B of Mah (2001) for details.

**Table B.2: Estimated Non-Fire Carbon Monoxide Poisoning Death Rates Associated with Generators, 1999–2018\***

Year	Estimate <sup>+</sup>	U.S. Population (thousands)	3-Year Average Mortality Rate per 10 Million Population
1999	7	279,040	
2000	19	282,172	0.54
2001	20	285,082	0.95
2002	42	287,804	1.29
2003	49	290,326	1.52
2004	41	293,046	2.02
2005	88	295,753	2.41
2006	85	298,593	2.69
2007	68	301,580	2.53
2008	76	304,375	2.28
2009	64	307,007	1.98
2010	42	309,338	1.83
2011	64	311,644	1.74
2012	57	313,993	1.88
2013	56	316,235	1.76
2014	54	318,857	2.03
2015	84	321,419	2.12
2016	66	323,128	2.54
2017	95	325,719	2.52
2018	84	326,838	2.85
2019	100	328,240	

\* Estimates are based on single source product incidents as multiple source incidents could be included in multiple categories.

+ Estimates in this table do not include multiple product-related deaths because a generator was not the sole product associated with the fatality.

Note 1: The 3-year average mortality rate is reported using the mid-year population estimates.

Note 2: Mortality rate changes from last year's report are due to changes in CPSC CO death estimates and changes in U.S. Census population estimates.

Table B.3 shows the CO poisoning mortality rates associated with all consumer products, excluding generators. The data indicate that, when generators are excluded, there does not appear to be a trend in the mortality rate for consumer product-related CO deaths. The 2000, 3-year annual average mortality rate was 3.60. The 2018, 3-year average mortality rate was 3.42, and the rate has risen each year since the 2016 low of 2.66.

**Table B.3: Estimated Non-Fire Carbon Monoxide Poisoning Death Rates Associated with Consumer Products (Excluding Generator-Related Deaths),\* 1999-2018**

Year	Estimate <sup>+</sup>	U.S. Population (thousands)	3-Year Average Mortality Rate per 10 Million Population
1999	95	279,040	
2000	117	282,172	3.60
2001	93	285,082	3.93
2002	126	287,804	3.65
2003	96	290,326	3.93
2004	120	293,046	3.48
2005	90	295,753	3.35
2006	87	298,593	3.07
2007	98	301,580	3.04
2008	90	304,375	2.86
2009	73	307,007	2.88
2010	102	309,338	2.87
2011	91	311,644	2.87
2012	75	313,993	2.66
2013	85	316,235	2.77
2014	103	318,857	2.79
2015	79	321,419	2.85
2016	92	323,128	2.66
2017	87	325,719	3.01
2018	114	326,838	3.42
2019	134	328,240	

\* Estimates are based on single source product incidents as multiple source incidents could be included in multiple categories.

+ Excludes estimates of deaths associated with a generator only.

Note 1: The 3-year average mortality rate is reported at the mid-year population estimates.

Note 2: Mortality rate changes from last year's report are due to changes in CPSC CO death estimates and changes in U.S. Census population estimates.

Table B.4 shows the 3-year average mortality rates of all engine-driven tools, including generators, through 2018. Although the average mortality rates for 2007 through 2011 have dropped slightly since the 2006 high (3.18), in 2018, the rate (3.28) increased to the highest rate since the 2007 rate of 2.93. The table shows that the 3-year average mortality rate has more than quadrupled from the rate in 2000 (0.71), to 2018 (3.28).

**Table B.4: Estimated Non-Fire Carbon Monoxide Poisoning Death Rates Associated with Generators and Other Engine-Driven Tools, 1999–2018\***

Year	Estimate <sup>+</sup>	U.S. Population (thousands)	3-Year Average Mortality Rate per 10 Million Population
1999	13	279,040	0.71
2000	26	282,172	
2001	21	285,082	1.16
2002	52	287,804	1.49
2003	56	290,326	1.88
2004	56	293,046	2.43
2005	102	295,753	2.95
2006	104	298,593	3.18
2007	79	301,580	2.93
2008	82	304,375	2.60
2009	76	307,007	2.32
2010	56	309,338	2.21
2011	73	311,644	2.06
2012	64	313,993	2.18
2013	68	316,235	2.06
2014	62	318,857	2.32
2015	92	321,419	2.43
2016	80	323,128	2.84
2017	104	325,719	2.91
2018	100	326,838	3.28
2019	118	328,240	

\* Estimates are based on single source product incidents as multiple source incidents could be included in multiple categories.

+ Estimates in this table do not include multiple product-related deaths because an EDT was not the sole product associated with the fatality. The one exception to this is the 2001 estimate that includes one estimated death associated with a generator and another EDT.

Note 1: The 3-year average mortality rate is reported at the mid-year population estimates.

Note 2: Mortality rate changes from last year's report are due to changes in CPSC CO death estimates and changes in U.S. Census population estimates.

Table B.5 shows the CO mortality rates associated with all consumer products, excluding generators and other engine-driven tools. The data indicate that the annual average, 3-year mortality rate decreased by about 13 percent for non-engine-driven tool consumer products (*i.e.*, excluding generators and other engine-driven tools), from the 2000 rate of 3.44, to the 2018 rate of 2.98. However, in the 14 years between 2005 and the current estimate for 2018, the non-EDT CO fatality rates has been relatively consistent, fluctuating in a narrow band between 2.37 and 2.98 per 10 million population.

**Table B.5: Estimated Non-Fire Carbon Monoxide Poisoning Death Rates Associated with Consumer Products (Excluding Generator- and Other Engine-Driven Tool-Related Deaths)\*, 1999–2019**

Year	Estimate <sup>+</sup>	U.S. Population (thousands)	3-Year Average Mortality Rate per 10 Million Population
1999	89	279,040	
2000	110	282,172	3.44
2001	92	285,082	3.72
2002	116	287,804	3.44
2003	89	290,326	3.56
2004	105	293,046	3.07
2005	76	295,753	2.81
2006	68	298,593	2.58
2007	87	301,580	2.64
2008	84	304,375	2.54
2009	61	307,007	2.53
2010	88	309,338	2.49
2011	82	311,644	2.55
2012	68	313,993	2.37
2013	73	316,235	2.49
2014	95	318,857	2.50
2015	71	321,419	2.55
2016	79	323,128	2.35
2017	78	325,719	2.62
2018	98	326,838	2.98
2019	116	328,240	

\* Estimates are based on single source product incidents as multiple source incidents could be included in multiple categories.

+ Excludes estimates of deaths associated with EDTs only. Multiproduct-associated incidents are included here because an EDT could not be identified as the only product involved. The one exception to this is the 2001 estimate, which excludes one estimated death associated with a generator and another EDT.

Note 1: The 3-year average mortality rate is reported at the mid-year population estimates.

Note 2: Mortality rate changes from last year's report are due to changes in CPSC CO death estimates and changes to U.S. Census estimates.

## **Summary of Tables B.1 – B.5**

When all consumer products are considered, there has been a 36 percent increase in the CO mortality rate from a 3-year average mortality rate of 4.34 in 2000, to 6.61 in 2018, as shown in Table B.1. Engine-driven tools and generators have had a substantial impact on the increase in the CO poisoning mortality rate involving consumer products. But, in recent years, non-generator-related CO fatalities have also been on the rise.

# Appendix C: Chi-Squared Test Results

## Age Group Test Result

Table 7 shows the estimated number of CO poisoning deaths categorized by victim age for the 11 most recent years of data (2009–2019). For the Chi-Square statistical analysis, the two younger groups (“Under 5” and “5–14”) were combined, due to their small, estimated averages. Chi-Square goodness-of-fit test results indicate a statistically significant difference between the proportion of CO victims in each age group from the general U.S. population. Each age group was analyzed separately, versus the expected proportion of the respective age group, based on U.S. population figures (assuming there was no age group effect on the CO poisoning fatality rate), to determine which age group proportions were significantly different from expectation. Binomial tests indicate that four of the five individual groups were found to be significantly different from what would be expected if there was no population group effect:

1. The “Under 15” group<sup>7</sup> was significantly lower ( $< 0.0001$ ).
2. The “15–24” group was significantly lower (0.0020).
3. The “45–64” group was significantly higher ( $< 0.0001$ ).
4. The “65 and over” group was also significantly higher ( $< 0.0001$ ).

## Gender Group Test Result

Table 8 presents the distribution of estimated CO deaths categorized by gender. For 2017–2019, the average percentage of male CO victims was also 76 percent, and the average percentage of female victims was 24 percent. By contrast, about 49 percent of the U.S. population is male, and 51 percent of the U.S. population is female.<sup>8</sup> The gender-related differences in CO Poisoning deaths were confirmed to be statistically significant ( $p$ -value  $< 0.0001$ ).

## Ethnicity/Race Group Test Result

Table 9 provides a summary of CO fatality victims characterized by race/ethnicity for the years 2008 through 2018. Estimates of the percentage of the U.S. population categorized into the various race/ethnicity groupings were based on single-race characterizations, as represented in the U.S. Census Bureau reports. Individuals reported as multi-race are included in the *Unknown/Other/Mixed* category.

Chi-square goodness-of-fit test results indicate a significant statistical difference ( $p$ -value = 0.0322) between the proportion of CO victims categorized by race/ethnicity from that of the general U.S. population. Each race/ethnicity group was analyzed separately, versus the expected proportion of the respective race/ethnicity group based on U.S. population figures,

---

<sup>7</sup> “Under 5” and “5–14” groups were combined “due to small sample sizes.

<sup>8</sup> Three-year average, 2016 to 2018, from July 2020 U.S. Census estimates of the U.S. population.

assuming there was no race/ethnicity group effect on the CO poisoning fatality rate. A Chi-Square statistical analysis was performed to determine which race/ethnicity group proportions were significantly greater than or less than the expectation. For the Chi-Square analysis, the three smaller groups (“Asian/Pacific,” “American Indian,” and “Unknown/Other/Mixed”) were combined, due to their relatively small proportion of the U.S. population. Binomial tests indicate that two race/ethnicity groups were statistically significantly different from the expected proportion based on the U.S. population. The observed proportion of Hispanic CO deaths was significantly lower (p-value of 0.0140) than the proportion of Hispanics in the U.S. population. Additionally, the observed proportion of White CO deaths was significantly higher (p-value = 0.0071) than the proportion of White Americans in the U.S. population. In previous years of this report, Black or African Americans demonstrated a significantly higher proportion of CO deaths than their proportion in the U.S. population and White Americans did not. This finding was similarly observed in the most recent previous (which included estimates up until 2018). Although this finding continues to be observed for the second year in a row, it is unclear if this is an anomaly or a pattern change.

## Appendix D: Regional Definitions

- 1) Northeast comprises New England and Middle Atlantic states.
  - a) New England: Maine, New Hampshire, Vermont, Massachusetts, Rhode Island, and Connecticut.
  - b) Middle Atlantic: New York, New Jersey, and Pennsylvania.
- 2) Midwest comprises East North Central and West North Central states.
  - a) East North Central: Ohio, Indiana, Illinois, Michigan, and Wisconsin.
  - b) West North Central: Minnesota, Iowa, Missouri, North Dakota, South Dakota, Nebraska, and Kansas.
- 3) South comprises South Atlantic, East South Central and West South-Central states.
  - a) South Atlantic: Delaware, Maryland, District of Columbia, Virginia, West Virginia, North Carolina, South Carolina, Georgia, and Florida.
  - b) East South Central: Kentucky, Tennessee, Alabama, and Mississippi.
  - c) West South Central: Arkansas, Louisiana, Oklahoma, and Texas.
- 4) West comprises Mountain and Pacific states.
  - a) Mountain: Montana, Idaho, Wyoming, Colorado, New Mexico, Arizona, Utah, and Nevada.
  - b) Pacific: Washington, Oregon, California, Alaska, and Hawaii

Source: U.S. Census Bureau 2012 Statistical Abstract <http://www.census.gov/compendia/statab/cats/population.html>

## References

Hnatov, Matthew V., Non-Fire Carbon Monoxide Deaths Associated with the Use of Consumer Products 2018 Annual Estimates. Washington, D.C.: U.S. Consumer Product Safety Commission. 2021.

Hnatov, Matthew V., *Fatal and Nonfatal Incidents Associated with Non-Fire Carbon Monoxide from Engine-Driven Generators and Other Engine-Driven Tools, 2011–2021*. Washington, D.C.: U.S. Consumer Product Safety Commission. 2022.

Inkster S.E. *Health Hazard Assessment of CO Poisoning Associated with Emissions from a Portable, 5.5 Kilowatt, Gasoline-Powered Generator*. Washington, D.C.: U.S. Consumer Product Safety Commission, 2004.

Long, K., Saltzman, L. (1995). Non-fire Carbon Monoxide Incidents: Morbidity and Mortality Related to the Use of Household Appliances. Washington, D.C.: U.S. Consumer Product Safety Commission.

Mah, J., Non-Fire Carbon Monoxide Deaths Associated with the Use of Consumer Products 1998 Annual Estimates. Washington, D.C.: U.S. Consumer Product Safety Commission. 2001.

U.S. Census Bureau, Population Division. *Annual Estimates of the Resident Population by Sex, Age, Race, and Hispanic Origin for the United States and States: April 1, 2010 to July 1, 2019. June 2020* <<https://www2.census.gov/programs-surveys/popest/tables/2010-2019/national/asrh/nc-est2019-sr11h.xlsx>>

U.S. Census Bureau, Population Division. *Annual Estimates of the Resident Population for the United States, Regions, States, and Puerto Rico: April 1, 2010 to July 1, 2019 (NST-EST2019-01)* <<https://www2.census.gov/programs-surveys/popest/tables/2010-2019/state/totals/nst-est2019-01.xlsx>>

U.S. Department of Agriculture, Economic Research Service. *2010 Rural-Urban Commuting Area Codes* <<http://www.ers.usda.gov/data-products/rural-urban-commuting-area-codes.aspx>>

Center for Rural Health, University of North Dakota School of Medicine and Health Sciences. ***Temporary Zip Ruca 3.10 File Access Page***

U.S. Office of Budget and Management. *OMB BULLETIN NO. 13-01: Revised Delineations of Metropolitan Statistical Areas, Micropolitan Statistical Areas, and Combined Statistical Areas, and Guidance on Uses of the Delineations of These Areas*. February 28, 2013